

Annexin VI Polyclonal Antibody

Catalog No :	YT0238
Reactivity :	Human;Rat;Mouse;
Applications :	WB;IHC;IF;ELISA
Target :	Annexin VI
Gene Name :	ANXA6
Protein Name :	Annexin A6
Human Gene Id :	309
Human Swiss Prot No :	P08133
Mouse Swiss Prot No :	P14824
Immunogen :	The antiserum was produced against synthesized peptide derived from human Annexin A6. AA range:1-50
Specificity :	Annexin VI Polyclonal Antibody detects endogenous levels of Annexin VI protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC: 1:100-300 ELISA: 1:20000. IF 1:100-300 Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	75kD


Background : Annexin VI belongs to a family of calcium-dependent membrane and phospholipid binding proteins. Several members of the annexin family have been implicated in membrane-related events along exocytotic and endocytotic pathways. The annexin VI gene is approximately 60 kbp long and contains 26 exons. It encodes a protein of about 68 kDa that consists of eight 68-amino acid repeats separated by linking sequences of variable lengths. It is highly similar to human annexins I and II sequences, each of which contain four such repeats. Annexin VI has been implicated in mediating the endosome aggregation and vesicle fusion in secreting epithelia during exocytosis. Alternatively spliced transcript variants have been described. [provided by RefSeq, Aug 2010],

Function : caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,domain:A pair of annexin repeats may form one binding site for calcium and phospholipid.,function:May associate with CD21. May regulate the release of Ca(2+) from intracellular stores.,induction:By EBV.,miscellaneous:Seems to bind one calcium ion with high affinity.,PTM:Phosphorylated in response to growth factor stimulation.,similarity:Belongs to the annexin family.,similarity:Contains 4 annexin repeats.,similarity:Contains 8 annexin repeats.,subcellular location:Identified by mass spectrometry in melanosome fractions from stage I to stage IV.,


Subcellular Location : Cytoplasm . Melanosome . Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

Expression : Adipocyte,Amygdala,Lung,Uterus,


Products Images




Immunofluorescence analysis of A549. 1,primary Antibody(red) was diluted at 1:200(4 °C overnight). 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 594 Secondary antibody was diluted at 1:1000(room temperature, 50min).3, Picture B: DAPI(blue) 10min.




Immunofluorescence analysis of HeLa cell. 1, Annexin VI Polyclonal Antibody (green) was diluted at 1:200 (4° overnight). (red) was diluted at 1:200 (4° overnight). 2, Goat Anti Rabbit Alexa Fluor 488 Catalog: RS3211 was diluted at 1:1000 (room temperature, 50min). Goat Anti Mouse Alexa Fluor 594 Catalog: RS3608 was diluted at 1:1000 (room temperature, 50min).




Western Blot analysis of various cells using Annexin VI Polyclonal Antibody diluted at 1:2000




Western Blot analysis of 293 cells using Annexin VI Polyclonal Antibody diluted at 1:2000



Immunofluorescence analysis of HeLa cells, using Annexin A6 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using Annexin A6 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from 293 cells, using Annexin A6 Antibody. The lane on the right is blocked with the synthesized peptide.