

## **CD45 Polyclonal Antibody**

Catalog No: YT0765

Reactivity: Human; Mouse; Rat

**Applications:** WB;IHC;IF;ELISA

Target: CD45

Fields: >>Cell adhesion molecules;>>T cell receptor signaling pathway;>>Fc gamma R-

mediated phagocytosis;>>Salmonella infection;>>Primary immunodeficiency

Gene Name: PTPRC

**Protein Name:** Receptor-type tyrosine-protein phosphatase C

P08575

P06800

Human Gene Id: 5788

**Human Swiss Prot** 

No:

**Mouse Swiss Prot** 

No:

Rat Gene ld: 24699

Rat Swiss Prot No: P04157

**Immunogen:** The antiserum was produced against synthesized peptide derived from human

CD45. AA range:981-1030

**Specificity:** CD45 Polyclonal Antibody detects endogenous levels of CD45 protein.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500 - 1:2000. IHC: 1:100-300 ELISA: 1:20000. IF 1:100-300 Not yet

tested in other applications.

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

1/3



chromatography using epitope-specific immunogen.

**Concentration**: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 147kD

Cell Pathway: Cell adhesion molecules (CAMs);T\_Cell\_Receptor;Fc gamma R-mediated

phagocytosis; Primary immunodeficiency;

**Background:** The protein encoded by this gene is a member of the protein tyrosine

phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced

transcripts variants of this gene, which enc

**Function:** alternative products:At least 8 isoforms are produced,catalytic activity:Protein

tyrosine phosphate + H(2)O = protein tyrosine + phosphate., disease: Defects in PTPRC are a cause of severe combined immunodeficiency autosomal recessive

T-cell-negative/B-cell-positive/NK-cell-positive (T(-)B(+)NK(+)SCID)

[MIM:608971]. SCID refers to a genetically and clinically heterogeneous group of rare congenital disorders characterized by impairment of both humoral and cell-mediated immunity, leukopenia, and low or absent antibody levels. Patients with SCID present in infancy with recurrent, persistent infections by opportunistic organisms. The common characteristic of all types of SCID is absence of T-cell-mediated cellular immunity due to a defect in T-cell development., disease: Genetic

variations in PTPRC are involved in multiple sclerosis susceptibility (MS)

[MIM:126200]. MS is a neurodegenerative dis

Subcellular Location:

Cell membrane; Single-pass type I membrane protein. Membrane raft.

Colocalized with DPP4 in membrane rafts. .


**Expression :** Isoform 1: Detected in thymocytes. Isoform 2: Detected in thymocytes. Isoform 3:

Detected in thymocytes. Isoform 4: Not detected in thymocytes. Isoform 5: Detected in thymocytes. Isoform 6: Not detected in thymocytes. Isoform 7:


Detected in thymocytes. Isoform 8: Not detected in thymocytes.




## **Products Images**



Immunofluorescence analysis of A549. 1,primary Antibody(red) was diluted at 1:200(4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 594 Secondary antibody was diluted at 1:1000(room temperature, 50min).3, Picture B: DAPI(blue) 10min.



Western Blot analysis of Jurkat cells using CD45 Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using CD45 Antibody. The picture on the right is blocked with the synthesized peptide.