

BRD3 Polyclonal Antibody

Catalog No :	YT0522
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	BRD3
Gene Name :	BRD3
Protein Name :	Bromodomain-containing protein 3
Human Gene Id :	8019
Human Swiss Prot No :	Q15059
Mouse Gene Id :	67382
Mouse Swiss Prot No :	Q8K2F0
Immunogen :	The antiserum was produced against synthesized peptide derived from human BRD3. AA range:611-660
Specificity :	BRD3 Polyclonal Antibody detects endogenous levels of BRD3 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 90kD


Background : This gene was identified based on its homology to the gene encoding the RING3 protein, a serine/threonine kinase. The gene localizes to 9q34, a region which contains several major histocompatibility complex (MHC) genes. The function of the encoded protein is not known. [provided by RefSeq, Jul 2008],

Function : disease:A chromosomal aberration involving BRD3 is found in a rare, aggressive, and lethal carcinoma arising in midline organs of young people. Translocation t(15;9)(q14;q34) with NUT which produces a BRD3-NUT fusion protein.,similarity:Contains 2 bromo domains.,tissue specificity:Ubiquitous.,


Subcellular Location : Nucleus . Detected on chromatin. .

Expression : Ubiquitous.


Products Images



Western Blot analysis of various cells using BRD3 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Western Blot analysis of A549 cells using BRD3 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).