

## **CaMKV Polyclonal Antibody**

YT0630 Catalog No:

Reactivity: Human; Mouse; Rat

**Applications:** WB;IHC;IF;ELISA

Target: CaMKV

Gene Name: **CAMKV** 

**Protein Name:** CaM kinase-like vesicle-associated protein

**Human Gene Id:** 79012

**Human Swiss Prot** 

No:

Q8NCB2

Mouse Gene Id:

235604

**Mouse Swiss Prot** 

No:

Q3UHL1

Rat Gene Id:

79011

Rat Swiss Prot No: Q63092

Immunogen:

The antiserum was produced against synthesized peptide derived from human

CAMK5. AA range:211-260

**Specificity:** 

CaMKV Polyclonal Antibody detects endogenous levels of CaMKV protein.

Formulation:

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source:

Polyclonal, Rabbit, IgG

**Dilution:** 

WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:10000. Not

yet tested in other applications.

**Purification:** 

The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



**Concentration:** 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 54kD

**Background:** cofactor:Calcium.,domain:The protein kinase domain is predicted to be

catalytically inactive.,function:Does not appear to have detectable kinase activity.,similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family.,similarity:Contains 1 protein kinase domain.,subcellular location:Predominantly observed in association with the plasma membrane of soma and in neurites, both axons and dendrites. May be associated with vesicular

structures., subunit: Interacts with calmodulin, in the presence of calcium.,

**Function:** cofactor:Calcium.,domain:The protein kinase domain is predicted to be

catalytically inactive.,function:Does not appear to have detectable kinase activity.,similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family.,similarity:Contains 1 protein kinase domain.,subcellular location:Predominantly observed in association with the plasma membrane of soma and in neurites, both axons and dendrites. May be associated with vesicular

structures., subunit: Interacts with calmodulin, in the presence of calcium.,


Subcellular Location:

Cell membrane; Peripheral membrane protein. Cytoplasmic vesicle membrane; Peripheral membrane protein. Predominantly observed in association with the plasma membrane of soma and in neurites, both axons and dendrites. May be


associated with vesicular structures (By similarity). .

**Expression:** Brain, Lung, Retinoblastoma, Teratocarcinoma,


## **Products Images**




Western Blot analysis of various cells using CaMKV Polyclonal Antibody



Immunofluorescence analysis of HeLa cells, using CAMK5 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using CAMK5 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HepG2 and HUVEC cells, using CAMK5 Antibody. The lane on the right is blocked with the synthesized peptide.