

CD160 Polyclonal Antibody

Catalog No :	YT0729
Reactivity :	Human;Mouse
Applications :	WB;IF;ELISA
Target :	CD160
Gene Name :	CD160
Protein Name :	CD160 antigen
Human Gene Id :	11126
Human Swiss Prot No :	O95971
Mouse Swiss Prot No :	O88875
Immunogen :	The antiserum was produced against synthesized peptide derived from human CD160. AA range:21-70
Specificity :	CD160 Polyclonal Antibody detects endogenous levels of CD160 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	17kD


Background : CD160 is an 27 kDa glycoprotein which was initially identified with the monoclonal antibody BY55. Its expression is tightly associated with peripheral blood NK cells and CD8 T lymphocytes with cytolytic effector activity. The cDNA sequence of CD160 predicts a cysteine-rich, glycosylphosphatidylinositol-anchored protein of 181 amino acids with a single Ig-like domain weakly homologous to KIR2DL4 molecule. CD160 is expressed at the cell surface as a tightly disulfide-linked multimer. RNA blot analysis revealed CD160 mRNAs of 1.5 and 1.6 kb whose expression was highly restricted to circulating NK and T cells, spleen and small intestine. Within NK cells CD160 is expressed by CD56dimCD16+ cells whereas among circulating T cells its expression is mainly restricted to TCRgd bearing cells and to TCRab+CD8brightCD95+CD56+CD28-CD27-cells. In tissues, CD160 is expressed on all intestinal intraepithelial lymphocytes. C

Function : function:Receptor showing broad specificity for both classical and non-classical MHC class I molecules.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,subunit:Homomultimer; disulfide-linked.,tissue specificity:Expressed in spleen, peripheral blood, and small intestine. Expression is restricted to functional NK and T cytotoxic lymphocytes.,


Subcellular Location : [CD160 antigen]: Cell membrane ; Lipid-anchor, GPI-anchor .; [CD160 antigen, soluble form]: Secreted. Released from the cell membrane by GPI cleavage. .

Expression : Expression is restricted to functional NK and cytotoxic T lymphocytes. Expressed in viral-specific effector memory and terminally differentiated effector memory CD8+ T cells. Expressed in memory and activated CD4+ T cell subsets (at protein level) (PubMed:9743336, PubMed:18193050, PubMed:11978774, PubMed:25255144). Expressed at high levels in intraepithelial lymphocytes (at protein level) (PubMed:9743336). Expressed in both alpha-beta and gamma-delta CD8+ T cell subsets (at protein level) (PubMed:9743336, PubMed:18193050, PubMed:11978774). Expressed in umbilical vein endothelial cells (at protein level) (PubMed:16809620). Expressed in monocytes and at lower levels in B cells (PubMed:23761635). Isoform 3: Expressed exclusively in activated NK cells (at protein level) (PubMed:19109136).

Products Images



Western Blot analysis of various cells using CD160 Polyclonal Antibody



Western blot analysis of lysates from Jurkat and COLO205 cells, using CD160 Antibody. The lane on the right is blocked with the synthesized peptide.