

DHRS4 Polyclonal Antibody

Catalog No :	YT1349
Reactivity :	Human;Rat;Mouse;
Applications :	WB;IHC;IF;ELISA
Target :	DHRS4
Fields :	>>Retinol metabolism;>>Metabolic pathways;>>Peroxisome
Gene Name :	DHRS4
Protein Name :	Dehydrogenase/reductase SDR family member 4
Human Gene Id :	10901
Human Swiss Prot No :	Q9BTZ2
Mouse Swiss Prot No :	Q99LB2
Immunogen :	The antiserum was produced against synthesized peptide derived from human DHRS4. AA range:191-240
Specificity :	DHRS4 Polyclonal Antibody detects endogenous levels of DHRS4 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 32-34kD

Cell Pathway : Retinol metabolism;


Background : catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,function:Reduces all-trans-retinal and 9-cis retinal. Can also catalyze the oxidation of all-trans-retinol with NADP as co-factor, but with much lower efficiency. Reduces alkyl phenyl ketones and alpha-dicarbonyl compounds with aromatic rings, such as pyrimidine-4-aldehyde, 3-benzoylpyridine, 4-benzoylpyridine, menadione and 4-hexanoylpyridine. Has no activity towards aliphatic aldehydes and ketones.,miscellaneous:Inhibited by kaempferol, quercetin, genistein and myristic acid.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,subcellular location:Isoform 1 is peroxisomal, while isoform 4 is not.,subunit:Homotetramer.,tissue specificity:Isoform 1 is predominantly expressed in normal cervix (at protein level). Isoform 4 is expressed in some neoplastic cervical tissues, but not in normal cervix (at protein level). Isoforms 5 and 6 are expressed in a few neoplastic cervical tissues.,

Function : catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,function:Reduces all-trans-retinal and 9-cis retinal. Can also catalyze the oxidation of all-trans-retinol with NADP as co-factor, but with much lower efficiency. Reduces alkyl phenyl ketones and alpha-dicarbonyl compounds with aromatic rings, such as pyrimidine-4-aldehyde, 3-benzoylpyridine, 4-benzoylpyridine, menadione and 4-hexanoylpyridine. Has no activity towards aliphatic aldehydes and ketones.,miscellaneous:Inhibited by kaempferol, quercetin, genistein and myristic acid.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,subcellular location:Isoform 1 is peroxisomal, while isoform 4 is not.,subunit:Homotetramer.,tissue specificity:Isoform 1 is predominantly expressed in normal cervix (at protein level). Isoform 4 is expressed in some neoplastic cervical tissues, but not in normal cervix (at protei


Subcellular Location : [Isoform 1]: Peroxisome . Isoform 4 is not peroxisomal. . ; [Isoform 7]: Nucleus .

Expression : [Isoform 1]: Predominantly expressed in normal cervix (at protein level). ; [Isoform 4]: Expressed in some neoplastic cervical tissues, but not in normal cervix (at protein level). ; [Isoform 5]: Expressed in a few neoplastic cervical tissues. ; [Isoform 6]: Expressed in a few neoplastic cervical tissues. ; [Isoform 8]: High expression in liver.


Products Images



Western Blot analysis of various cells using DHRS4 Polyclonal Antibody diluted at 1:2000



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using DHRS4 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HepG2 cells, using DHRS4 Antibody. The lane on the right is blocked with the synthesized peptide.