

DNAM-1 Polyclonal Antibody

| | |
|------------------------------|---|
| Catalog No : | YT1381 |
| Reactivity : | Human;Mouse;Monkey |
| Applications : | WB;IHC;IF;ELISA |
| Target : | DNAM-1 |
| Fields : | >>Cell adhesion molecules |
| Gene Name : | CD226 |
| Protein Name : | CD226 antigen |
| Human Gene Id : | 10666 |
| Human Swiss Prot No : | Q15762 |
| Mouse Gene Id : | 225825 |
| Mouse Swiss Prot No : | Q8K4F0 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human CD226/DNAM-1. AA range:287-336 |
| Specificity : | DNAM-1 Polyclonal Antibody detects endogenous levels of DNAM-1 protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 38kD

Cell Pathway : Cell adhesion molecules (CAMs);


Background : This gene encodes a glycoprotein expressed on the surface of NK cells, platelets, monocytes and a subset of T cells. It is a member of the Ig-superfamily containing 2 Ig-like domains of the V-set. The protein mediates cellular adhesion of platelets and megakaryocytic cells to vascular endothelial cells. The protein also plays a role in megakaryocytic cell maturation. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2015],

Function : function:Receptor involved in intercellular adhesion, lymphocyte signaling, cytotoxicity and lymphokine secretion mediated by cytotoxic T-lymphocyte (CTL) and NK cell.,PTM:Phosphorylated.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Interacts with PVR and PVRL2.,tissue specificity:Expressed by peripheral blood T-lymphocytes.,


Subcellular Location : Cell membrane ; Single-pass type I membrane protein .

Expression : Expressed by peripheral blood T-lymphocytes.


Products Images




Western Blot analysis of COS7 cells using DNAM-1 Polyclonal Antibody diluted at 1:500




Western Blot analysis of mouse-brain cells using DNAM-1 Polyclonal Antibody diluted at 1:500




Western Blot analysis of AD293 using DNAM-1 Polyclonal Antibody diluted at 1:500




Immunofluorescence analysis of HeLa cells, using CD226/DNAM-1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from COS7 and K562 cells, treated with PMA 125ng/ml 30', using CD226/DNAM-1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HUVEC cells using CD226/DNAM-1 antibody.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4° overnight). 2, Tris-EDTA, pH 9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).