

Fe65L Polyclonal Antibody

Catalog No :	YT1688
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	Fe65L
Gene Name :	APBB2
Protein Name :	Amyloid beta A4 precursor protein-binding family B member 2
Human Gene Id :	323
Human Swiss Prot No :	Q92870
Mouse Gene Id :	11787
Mouse Swiss Prot No :	Q9DBR4
Immunogen :	The antiserum was produced against synthesized peptide derived from human APBB2. AA range:471-520
Specificity :	Fe65L Polyclonal Antibody detects endogenous levels of Fe65L protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 83kD


Background : amyloid beta precursor protein binding family B member 2(APBB2) Homo sapiens The protein encoded by this gene interacts with the cytoplasmic domains of amyloid beta (A4) precursor protein and amyloid beta (A4) precursor-like protein 2. This protein contains two phosphotyrosine binding (PTB) domains, which are thought to function in signal transduction. Polymorphisms in this gene have been associated with Alzheimer's disease. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2009],

Function : function:May modulate the internalization of beta-amyloid precursor protein.,similarity:Contains 1 WW domain.,similarity:Contains 2 PID domains.,subunit:Binds to the intracellular domain of the beta-amyloid precursor protein.,


Subcellular Location : Endoplasmic reticulum . Golgi apparatus . Early endosome .

Expression : Widely expressed.


Products Images



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using APBB2 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from LOVO cells, using APBB2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using APBB2 antibody.