

EphA3 Polyclonal Antibody

| | |
|------------------------------|---|
| Catalog No : | YT1578 |
| Reactivity : | Human;Mouse;Rat |
| Applications : | WB;ELISA |
| Target : | EphA3 |
| Fields : | >>Axon guidance |
| Gene Name : | EPHA3 |
| Protein Name : | Ephrin type-A receptor 3 |
| Human Gene Id : | 2042 |
| Human Swiss Prot No : | P29320 |
| Mouse Swiss Prot No : | P29319 |
| Rat Gene Id : | 29210 |
| Rat Swiss Prot No : | O08680 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human EPHA3. AA range:831-880 |
| Specificity : | EphA3 Polyclonal Antibody detects endogenous levels of EphA3 protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |

| | |
|-------------------------------|---|
| Concentration : | 1 mg/ml |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |
| Observed Band : | 120kD |
| Cell Pathway : | Axon guidance; |
| Background : | <p>This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. This gene encodes a protein that binds ephrin-A ligands. Two alternatively spliced transcript variants have been described for this gene. [provided by RefSeq, Jul 2008],</p> |
| Function : | <p>catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,disease:Defects in EPHA3 may be a cause of colorectal cancer (CRC) [MIM:114500].,function:Receptor for members of the ephrin-A family. Binds to ephrin-A2, -A3, -A4 and -A5. Could play a role in lymphoid function.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. Ephrin receptor subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,similarity:Contains 2 fibronectin type-III domains.,tissue specificity:Widely expressed. Highest level in placenta.,</p> |
| Subcellular Location : | [Isoform 1]: Cell membrane ; Single-pass type I membrane protein .; [Isoform 2]: Secreted . |
| Expression : | Widely expressed. Highest level in placenta. |

Products Images

Western Blot analysis of 293T-UV cells using EphA3 Polyclonal Antibody diluted at 1:500

