

DOR-1 Polyclonal Antibody

Catalog No: YT1403

Reactivity: Human; Mouse; Rat; Monkey

Applications: WB;IHC;IF;ELISA

Target: DOR-1

Fields: >>cGMP-PKG signaling pathway;>>Sphingolipid signaling

pathway;>>Neuroactive ligand-receptor interaction

Gene Name: OPRD1

Protein Name : Delta-type opioid receptor

P41143

P32300

Human Gene Id: 4985

Human Swiss Prot

No:

Mouse Gene Id: 18386

Mouse Swiss Prot

No:

Rat Gene Id: 24613

Rat Swiss Prot No: P33533

Immunogen: The antiserum was produced against synthesized peptide derived from human

Opioid Receptor-delta. AA range:323-372

Specificity: DOR-1 Polyclonal Antibody detects endogenous levels of DOR-1 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution : WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200

1/3



Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 40kD

Cell Pathway: Neuroactive ligand-receptor interaction;

Background: function:Inhibits neurotransmitter release by reducing calcium ion currents and

increasing potassium ion conductance. Highly stereoselective. receptor for enkephalins.,online information:Delta opioid receptor entry,similarity:Belongs to the G-protein coupled receptor 1 family.,subunit:Interacts with GPRASP1.,

and a protein oddpiod rosopter i lanning, jodd annanteracte with an interest

Function: function:Inhibits neurotransmitter release by reducing calcium ion currents and

increasing potassium ion conductance. Highly stereoselective. receptor for enkephalins.,online information:Delta opioid receptor entry,similarity:Belongs to

the G-protein coupled receptor 1 family., subunit: Interacts with GPRASP1.,

Subcellular Location:


Cell membrane; Multi-pass membrane protein.

Expression: Detected in oocytes (at protein level). Detected in brain cortex, hypothalamus,


hippocampus and olfactory bulb. Detected in oocytes.

Products Images





Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Opioid Receptor-delta Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HUVEC and COLO205 cells, using Opioid Receptor-delta Antibody. The lane on the right is blocked with the synthesized peptide.