

## EMR1 Polyclonal Antibody

| Catalog No :        | YT1544  |
|---------------------|---|
| Reactivity :        | Human;Rat;Mouse;  |
| Applications :      | WB;ELISA  |
| Target :            | EMR1  |
| Gene Name :         | EMR1  |
| Protein Name :      | EGF-like module-containing mucin-like hormone receptor-like 1   |
| Human Gene Id :     | 2015  |
| Human Swiss Prot    | Q14246  |
| Mouse Swiss Prot    | Q61549  |
| Immunogen :         | The antiserum was produced against synthesized peptide derived from human EMR1. AA range:818-867                          |
| Specificity :       | EMR1 Polyclonal Antibody detects endogenous levels of EMR1 protein.   |
| Formulation :       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source :            | Polyclonal, Rabbit,IgG  |
| Dilution :          | WB 1:500 - 1:2000. ELISA: 1:5000. Not yet tested in other applications.   |
| Purification :      | The antibody was affinity-purified from rabbit antiserum by affinity-<br>chromatography using epitope-specific immunogen. |
| Concentration :     | 1 mg/ml   |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C)  |
| Observed Band :     | 98kD  |



| Background :                              | This gene encodes a protein that has a domain resembling seven<br>transmembrane G protein-coupled hormone receptors (7TM receptors) at its C-<br>terminus. The N-terminus of the encoded protein has six EGF-like modules,<br>separated from the transmembrane segments by a serine/threonine-rich domain,<br>a feature reminiscent of mucin-like, single-span, integral membrane glycoproteins<br>with adhesive properties. Multiple alternatively spliced transcript variants<br>encoding different isoforms have been found for this gene. [provided by RefSeq,<br>Jan 2012], |
|---|--|
| Function :                                | alternative products:Comment=Additional isoforms seem to exist,function:Could<br>be involved in cell-cell interactions.,similarity:Belongs to the G-protein coupled<br>receptor 2 family. LN-TM7 subfamily.,similarity:Contains 1 GPS<br>domain.,similarity:Contains 6 EGF-like domains.,tissue specificity:Wide<br>expression; increased levels in peripheral blood mononuclear cells.,   |
| Subcellular<br>Location :<br>Expression : | Cell membrane ; Multi-pass membrane protein .<br>Expression is restricted to eosinophils.  |





