

## GPR133 Polyclonal Antibody

<b>Catalog No :</b>	YT1969
<b>Reactivity :</b>	Human;Monkey
<b>Applications :</b>	WB;IF;ELISA
<b>Target :</b>	GPR133
<b>Gene Name :</b>	GPR133
<b>Protein Name :</b>	Probable G-protein coupled receptor 133
<b>Human Gene Id :</b>	283383
<b>Human Swiss Prot No :</b>	Q6QNK2
<b>Mouse Swiss Prot No :</b>	Q80T32
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human GPR133. AA range:461-510
<b>Specificity :</b>	GPR133 Polyclonal Antibody detects endogenous levels of GPR133 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:5000. Not yet tested in other applications.
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	96kD


**Background :** The adhesion G-protein-coupled receptors (GPCRs), including GPR133, are membrane-bound proteins with long N termini containing multiple domains. GPCRs, or GPRs, contain 7 transmembrane domains and transduce extracellular signals through heterotrimeric G proteins (summary by Bjarnadottir et al., 2004 [PubMed 15203201]).[supplied by OMIM, Nov 2010],

**Function :** function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 2 family. LN-TM7 subfamily.,similarity:Contains 1 GPS domain.,


**Subcellular Location :** Cell membrane ; Multi-pass membrane protein .

**Expression :** Up-regulated in CD133(+) cell population of glioblastoma.


## Products Images



Immunofluorescence analysis of HeLa cells, using GPR133 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from COS7 cells, using GPR133 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from K562 cells using GPR133 antibody.