

GPR20 Polyclonal Antibody

Catalog No: YT2008

Reactivity: Human; Rat; Mouse;

Applications: WB;IHC;IF;ELISA

Target: GPR20

Gene Name: GPR20

Protein Name: G-protein coupled receptor 20

Q99678

Q8BYC4

Human Gene ld: 2843

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

GPR20. AA range:291-340

Specificity: GPR20 Polyclonal Antibody detects endogenous levels of GPR20 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source : Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:5000. Not

yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 42kD



Background: function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1

family.,tissue specificity:Liver and the brain regions thalamus, putamen, and

caudate, but not in frontal cortex, pons and hypothalamus.,

Function: function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1

family.,tissue specificity:Liver and the brain regions thalamus, putamen, and

caudate, but not in frontal cortex, pons and hypothalamus.,

Subcellular Location:


Cell membrane; Multi-pass membrane protein.

Expression: Ubiquitous with highest levels in intestinal tissues. In the brain detected in


thalamus, putamen, and caudate, but not in frontal cortex, pons and

hypothalamus.


Products Images



Immunofluorescence analysis of A549 cells, using GPR20 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using GPR20 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from LOVO cells, using GPR20 Antibody. The lane on the right is blocked with the synthesized peptide.