

GDF-15 Polyclonal Antibody

Catalog No :	YT1888
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA
Target :	GDF-15
Fields :	>>Cytokine-cytokine receptor interaction
Gene Name :	GDF15
Protein Name :	Growth/differentiation factor 15
Human Gene Id :	9518
Human Swiss Prot No :	Q99988
Mouse Swiss Prot No :	Q9Z0J7
Immunogen :	The antiserum was produced against synthesized peptide derived from human GDF15. AA range:31-80
Specificity :	GDF-15 Polyclonal Antibody detects endogenous levels of GDF-15 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 37kD


Background : growth differentiation factor 15(GDF15) Homo sapiens This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. The protein is expressed in a broad range of cell types, acts as a pleiotropic cytokine and is involved in the stress response program of cells after cellular injury. Increased protein levels are associated with disease states such as tissue hypoxia, inflammation, acute injury and oxidative stress. [provided by RefSeq, Aug 2016],

Function : similarity:Belongs to the TGF-beta family.,subunit:Homodimer; disulfide-linked.,tissue specificity:Highly expressed in placenta, with lower levels in prostate and colon and some expression in kidney.,


Subcellular Location : Secreted .

Expression : Highly expressed in placenta, with lower levels in prostate and colon and some expression in kidney (PubMed:9348093). Detected in plasma (at protein level) (PubMed:28572090, PubMed:29046435).


Products Images



Western Blot analysis of various cells using GDF-15 Polyclonal Antibody diluted at 1:1000



Western blot analysis of lysates from 293 and HepG2 cells, using GDF15 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using GDF15 antibody.