

HSP105 Polyclonal Antibody

Catalog No :	YT2249
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	HSP105
Fields :	>>Protein processing in endoplasmic reticulum
Gene Name :	HSPH1
Protein Name :	Heat shock protein 105 kDa
Human Gene Id :	10808
Human Swiss Prot No :	Q92598
Mouse Gene Id :	15505
Mouse Swiss Prot No :	Q61699
Immunogen :	The antiserum was produced against synthesized peptide derived from human HSP105. AA range:791-840
Specificity :	HSP105 Polyclonal Antibody detects endogenous levels of HSP105 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 100kD


Background : function:Prevents the aggregation of denatured proteins in cells under severe stress, on which the ATP levels decrease markedly. Inhibits HSPA8/HSC70 ATPase and chaperone activities.,PTM:Phosphorylation on Ser-509 may be important for regulation of the HSPA8/HSC70 chaperone activity.,similarity:Belongs to the heat shock protein 70 family.,subunit:Interacts with HSPA8/HSC70.,tissue specificity:Highly expressed in testis. Present at lower levels in most brain regions, except cerebellum. Overexpressed in cancer cells.,

Function : function:Prevents the aggregation of denatured proteins in cells under severe stress, on which the ATP levels decrease markedly. Inhibits HSPA8/HSC70 ATPase and chaperone activities.,PTM:Phosphorylation on Ser-509 may be important for regulation of the HSPA8/HSC70 chaperone activity.,similarity:Belongs to the heat shock protein 70 family.,subunit:Interacts with HSPA8/HSC70.,tissue specificity:Highly expressed in testis. Present at lower levels in most brain regions, except cerebellum. Overexpressed in cancer cells.,


Subcellular Location : Cytoplasm .

Expression : Highly expressed in testis. Present at lower levels in most brain regions, except cerebellum. Overexpressed in cancer cells.


Products Images




Western Blot analysis of various cells using HSP105 Polyclonal Antibody



Immunofluorescence analysis of HeLa cells, using HSP105 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using HSP105 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa cells, using HSP105 Antibody. The lane on the right is blocked with the synthesized peptide.