

Phospho GSK3β(Ser9) Mouse mAb(6H6)

Catalog No: YP1886

Reactivity: Human; Mouse; Rat

Applications: IHC;WB

Target: GSK3β

Fields: >>EGFR tyrosine kinase inhibitor resistance;>>ErbB signaling

pathway;>>Chemokine signaling pathway;>>Cell cycle;>>mTOR signaling pathway;>>PI3K-Akt signaling pathway;>>Wnt signaling pathway;>>Hedgehog signaling pathway;>>Axon guidance;>>Hippo signaling pathway;>>Focal adhesion;>>Signaling pathways regulating pluripotency of stem cells;>>IL-17 signaling pathway;>>T cell receptor signaling pathway;>>B cell receptor signaling pathway;>>Neurotrophin signaling pathway;>>Dopaminergic synapse;>>Insulin signaling pathway;>>Melanogenesis;>>Prolactin signaling pathway;>>Thyroid hormone signaling pathway;>>Insulin resistance;>>Non-alcoholic fatty liver disease;>>Cushing syndrome;>>Growth hormone synthesis, secretion and

action;>>Alcoholic liver disease;>>Alzheimer disease;>>Prion

disease;>>Pathways of neurodegeneration - multiple

diseases;>>Shigellosis;>>Yersinia infection;>>Hepatitis C;>>Measles;>>Human cytomegalovirus infection;>>Human papillomavirus infection;>>Kaposi sarcoma-

associated herpes

Gene Name: GSK3B

Protein Name : Phospho-GSK3β(Ser9)

Q9WV60

Human Gene Id: 2932

Human Swiss Prot P49841

No:

Mouse Gene Id: 56637

Mouse Swiss Prot

No:

Rat Gene Id: 84027

Rat Swiss Prot No: P18266

1/3



Immunogen : Synthesized peptide derived from human Phospho-GSK3β(Ser9)

Specificity: This antibody detects endogenous levels of Phospho-GSK3β(Ser9) at Human,

Mouse,Rat

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Monoclonal, Mouse

Dilution : IHC1:50-200 ,WB 1:1000-2000

Purification: The antibody was affinity-purified from mouse ascites by affinity-

chromatography using specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 46kD

Background: The protein encoded by this gene is a serine-threonine kinase, belonging to the

glycogen synthase kinase subfamily. It is involved in energy metabolism, neuronal cell development, and body pattern formation. Polymorphisms in this gene have been implicated in modifying risk of Parkinson disease, and studies in mice show that overexpression of this gene may be relevant to the pathogenesis of Alzheimer disease. Alternatively spliced transcript variants encoding different isoforms have

been found for this gene.[provided by RefSeq, Sep 2009],

Function : catalytic activity:ATP + [tau protein] = ADP + [tau protein] phosphate.,enzyme

regulation:Inhibited when phosphorylated by AKT1.,function:Participates in the Wnt signaling pathway. Implicated in the hormonal control of several regulatory proteins including glycogen synthase, MYB and the transcription factor JUN. Phosphorylates JUN at sites proximal to its DNA-binding domain, thereby reducing its affinity for DNA. Phosphorylates MUC1 in breast cancer cells, and

decreases the interaction of MUC1 with CTNNB1/beta-

catenin.,PTM:Phosphorylated by AKT1 and ILK1.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. GSK-3 subfamily.,similarity:Contains 1 protein kinase domain.,subunit:Monomer (By similarity). Interacts with CABYR, MUC1,

NIN and PRUNE., tissue specificity: Expressed in testis, thymus, prostate

Subcellular Location:


Cytoplasm . Nucleus . Cell membrane . The phosphorylated form shows localization to cytoplasm and cell membrane (PubMed:20937854). The MEMO1-RHOA-DIAPH1 signaling pathway controls localization of the phosphorylated form to the cell membrane (PubMed:20937854). .




Expression:

Expressed in testis, thymus, prostate and ovary and weakly expressed in lung, brain and kidney. Colocalizes with EIF2AK2/PKR and TAU in the Alzheimer disease (AD) brain.

Products Images



Immunohistochemical analysis of paraffin-embedded Human Lung Carcinoma Tissue using GSK3 β Mouse mAb diluted at 1:200.



Western blot analysis of 1) Hela Cell Lysate, 2) Mouse Heart Tissue Lysate, 3)Rat Heart Tissue Lysate using GSK 3β Mouse mAb diluted at 1:1000.