


MCT4 Polyclonal Antibody


Catalog No :	YT2685
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	MCT4
Fields :	>>Central carbon metabolism in cancer
Gene Name :	SLC16A3
Protein Name :	Monocarboxylate transporter 4
Human Gene Id :	9123
Human Swiss Prot No :	O15427
Mouse Gene Id :	80879
Mouse Swiss Prot No :	P57787
Rat Gene Id :	80878
Rat Swiss Prot No :	O35910
Immunogen :	The antiserum was produced against synthesized peptide derived from human MOT4. AA range:233-282
Specificity :	MCT4 Polyclonal Antibody detects endogenous levels of MCT4 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications.

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	49kD
Background :	Lactic acid and pyruvate transport across plasma membranes is catalyzed by members of the proton-linked monocarboxylate transporter (MCT) family, which has been designated solute carrier family-16. Each MCT appears to have slightly different substrate and inhibitor specificities and transport kinetics, which are related to the metabolic requirements of the tissues in which it is found. The MCTs, which include MCT1 (SLC16A1; MIM 600682) and MCT2 (SLC16A7; MIM 603654), are characterized by 12 predicted transmembrane domains (Price et al., 1998 [PubMed 9425115]).[supplied by OMIM, Mar 2008],
Function :	function:Proton-linked monocarboxylate transporter. Catalyzes the rapid transport across the plasma membrane of many monocarboxylates such as lactate, pyruvate, branched-chain oxo acids derived from leucine, valine and isoleucine, and the ketone bodies acetoacetate, beta-hydroxybutyrate and acetate.,similarity:Belongs to the major facilitator superfamily. Monocarboxylate porter (TC 2.A.1.13) family.,tissue specificity:Highly expressed in skeletal muscle.,
Subcellular Location :	Cell membrane; Multi-pass membrane protein.
Expression :	Highly expressed in skeletal muscle.


Products Images



Western Blot analysis of various cells using MCT4 Polyclonal Antibody diluted at 1:500



Western Blot analysis of PC-3 cells using MCT4 Polyclonal Antibody diluted at 1:500



Western blot analysis of lysates from LOVO cells, using MOT4 Antibody. The lane on the right is blocked with the synthesized peptide.