

### **Olfactory receptor 2B8 Polyclonal Antibody**

Catalog No: YT3300

**Reactivity:** Human; Rat; Mouse;

**Applications:** WB;IF;ELISA

Target: Olfactory receptor 2B8

Gene Name: OR2B8P

**Protein Name:** Putative olfactory receptor 2B8

P59922

Human Gene Id: 65944

**Human Swiss Prot** 

No:

**Immunogen:** The antiserum was produced against synthesized peptide derived from human

OR2B8. AA range:61-110

**Specificity:** Olfactory receptor 2B8 Polyclonal Antibody detects endogenous levels of

Olfactory receptor 2B8 protein.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**Source :** Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other

applications.

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 38kD

1/2



#### **Background:**

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],


#### **Function:**

caution:Could be the product of a pseudogene.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,


# Subcellular Location:

Cell membrane; Multi-pass membrane protein.

## **Products Images**



Western blot analysis of lysates from HeLa cells, using OR2B8 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using OR2B8 antibody.