

Olfactory receptor 52A4 Polyclonal Antibody

Catalog No: YT3374

Reactivity: Human; Rat; Mouse;

Applications: WB;IF;ELISA

Target: Olfactory receptor 52A4

Gene Name: OR52A4

Protein Name: Putative olfactory receptor 52A4

A6NMU1

Human Gene Id: 390053

Human Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

OR52A4. AA range:211-260

Specificity: Olfactory receptor 52A4 Polyclonal Antibody detects endogenous levels of

Olfactory receptor 52A4 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other

applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 35kD

1/2



Background:

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. Although originally considered to be a functional olfactory receptor, this family member is now considered to be pseudogene due to the presence of a C-terminal frameshift compared to other family membe


Function:

caution: Could be the product of a pseudogene. Has a truncated and frameshifted C-terminal region., similarity: Belongs to the G-protein coupled receptor 1 family.,


Subcellular Location:

Cell membrane; Multi-pass membrane protein.

Products Images



Western blot analysis of lysates from MCF-7 cells, using OR52A4 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from K562 cells using OR52A4 antibody.