

Olfactory receptor 52N4 Polyclonal Antibody

| | |
|------------------------------|---|
| Catalog No : | YT3385 |
| Reactivity : | Human;Rat;Mouse; |
| Applications : | WB;IF;ELISA |
| Target : | Olfactory receptor 52N4 |
| Fields : | >>Olfactory transduction |
| Gene Name : | OR52N4 |
| Protein Name : | Olfactory receptor 52N4 |
| Human Gene Id : | 390072 |
| Human Swiss Prot No : | Q8NGI2 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human OR52N4. AA range:272-321 |
| Specificity : | Olfactory receptor 52N4 Polyclonal Antibody detects endogenous levels of Olfactory receptor 52N4 protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |

Observed Band : 36kD


Cell Pathway : Olfactory transduction;

Background : Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],


Function : function:Odorant receptor .,polymorphism:A stop codon at position Arg-172 in the gene coding for this protein is responsible for functional diversity thus producing a pseudogene. The stop codon is more frequent in non-Africans than in African-Americans.,similarity:Belongs to the G-protein coupled receptor 1 family.,

Subcellular Location : Cell membrane; Multi-pass membrane protein.


Products Images



Western Blot analysis of various cells using Olfactory receptor 52N4 Polyclonal Antibody



Western blot analysis of lysates from HeLa cells, using OR52N4 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using OR52N4 antibody.