

## **Olfactory receptor 5D16 Polyclonal Antibody**

Catalog No: YT3401

**Reactivity:** Human; Rat; Mouse;

**Applications:** WB;ELISA

Target: Olfactory receptor 5D16

**Fields:** >>Olfactory transduction

Gene Name: OR5D16

**Protein Name:** Olfactory receptor 5D16

Q8NGK9

Human Gene Id: 390144

**Human Swiss Prot** 

No:

**Immunogen:** The antiserum was produced against synthesized peptide derived from human

OR5D16. AA range:201-250

**Specificity:** Olfactory receptor 5D16 Polyclonal Antibody detects endogenous levels of

Olfactory receptor 5D16 protein.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500 - 1:2000. ELISA: 1:5000. Not yet tested in other applications.

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 38kD

**Cell Pathway:** Olfactory transduction;

**Background :** Olfactory receptors interact with odorant molecules in the nose, to initiate a

neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR)

arising from single coding-exon genes. Olfactory receptors share a

7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by

RefSeq, Jul 2008],


Function: function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor

1 family.,

Subcellular Location:

Cell membrane; Multi-pass membrane protein.

## **Products Images**



Western blot analysis of lysates from MCF-7 cells, using OR5D16 Antibody. The lane on the right is blocked with the synthesized peptide.