

Pepsin C Polyclonal Antibody

Catalog No: YT3661

Reactivity: Human; Rat; Mouse;

Applications: WB;IHC;IF;ELISA

Target: Pepsin C

Gene Name: PGC

Protein Name: Gastricsin

Human Gene ld: 5225

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Immunogen:

Q9D7R7

P20142

The antiserum was produced against synthesized peptide derived from human

Pepsin C. AA range:257-306

Specificity: Pepsin C Polyclonal Antibody detects endogenous levels of Pepsin C protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source : Polyclonal, Rabbit, IgG

Dilution : WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 42kD

1/2



Background:

This gene encodes an aspartic proteinase that belongs to the peptidase family A1. The encoded protein is a digestive enzyme that is produced in the stomach and constitutes a major component of the gastric mucosa. This protein is also secreted into the serum. This protein is synthesized as an inactive zymogen that includes a highly basic prosegment. This enzyme is converted into its active mature form at low pH by sequential cleavage of the prosegment that is carried out by the enzyme itself. Polymorphisms in this gene are associated with susceptibility to gastric cancers. Serum levels of this enzyme are used as a biomarker for certain gastric diseases including Helicobacter pylori related gastritis. Alternate splicing results in multiple transcript variants. A pseudogene of this gene is found on chromosome 1. [provided by RefSeq, Oct 2009],

Function:

catalytic activity:More restricted specificity than pepsin A, but shows preferential cleavage at Tyr-|-Xaa bonds. High activity on hemoglobin.,function:Hydrolyzes a variety of proteins.,similarity:Belongs to the peptidase A1 family.,


Subcellular Location:

Secreted.


Expression:

Brain, Liver, Placenta, Stomach,

Products Images



Immunohistochemistry analysis of Pepsin C antibody in paraffinembedded human lung carcinoma tissue.



Western blot analysis of lysate from Jurkat cells, using Pepsin C antibody.