


## PDK2 Polyclonal Antibody


<b>Catalog No :</b>	YT3646
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	PDK2
<b>Fields :</b>	>>Diabetic cardiomyopathy
<b>Gene Name :</b>	PDK2
<b>Protein Name :</b>	[Pyruvate dehydrogenase [lipoamide]] kinase isozyme 2 mitochondrial
<b>Human Gene Id :</b>	5164
<b>Human Swiss Prot No :</b>	Q15119
<b>Mouse Gene Id :</b>	18604
<b>Mouse Swiss Prot No :</b>	Q9JK42
<b>Rat Gene Id :</b>	81530
<b>Rat Swiss Prot No :</b>	Q64536
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human PDK2. AA range:260-309
<b>Specificity :</b>	PDK2 Polyclonal Antibody detects endogenous levels of PDK2 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200

<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	46kD
<b>Background :</b>	This gene encodes a member of the pyruvate dehydrogenase kinase family. The encoded protein phosphorylates pyruvate dehydrogenase, down-regulating the activity of the mitochondrial pyruvate dehydrogenase complex. Overexpression of this gene may play a role in both cancer and diabetes. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2010],
<b>Function :</b>	catalytic activity:ATP + [pyruvate dehydrogenase (acetyl-transferring)] = ADP + [pyruvate dehydrogenase (acetyl-transferring)] phosphate.,function:Inhibits the mitochondrial pyruvate dehydrogenase complex by phosphorylation of the E1 alpha subunit, thus contributing to the regulation of glucose metabolism.,similarity:Belongs to the PDK/BCKDK protein kinase family.,similarity:Contains 1 histidine kinase domain.,tissue specificity:Expressed in many tissues, with the highest level in heart and skeletal muscle, intermediate levels in brain, kidney, pancreas and liver, and low levels in placenta and lung.,
<b>Subcellular Location :</b>	Mitochondrion matrix.
<b>Expression :</b>	Expressed in many tissues, with the highest level in heart and skeletal muscle, intermediate levels in brain, kidney, pancreas and liver, and low levels in placenta and lung.


## Products Images




Western Blot analysis of various cells using PDK2 Polyclonal Antibody diluted at 1:500



Western blot analysis of lysates from HepG2 cells, using PDK2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using PDK2 antibody.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).