


**PFAAP5 Polyclonal Antibody**


<b>Catalog No :</b>	YT3679
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB;ELISA;IHC
<b>Target :</b>	PFAAP5
<b>Gene Name :</b>	N4BP2L2
<b>Protein Name :</b>	NEDD4-binding protein 2-like 2
<b>Human Gene Id :</b>	10443
<b>Human Swiss Prot No :</b>	Q92802
<b>Mouse Gene Id :</b>	381695
<b>Mouse Swiss Prot No :</b>	Q8JZS6
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human N4BP2L2. AA range:358-407
<b>Specificity :</b>	PFAAP5 Polyclonal Antibody detects endogenous levels of PFAAP5 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

<b>Observed Band :</b>	<u>67kD</u>
<b>Background :</b>	<u>PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,</u>
<b>Function :</b>	<u>PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,</u>
<b>Subcellular Location :</b>	<u>nucleus,transcriptional repressor complex,extracellular exosome,</u>
<b>Expression :</b>	<u>Colon,Thalamus,</u>


## Products Images




Western Blot analysis of LOVO cells using PFAAP5 Polyclonal Antibody



Western blot analysis of lysates from LOVO cells, using N4BP2L2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from K562 cells using N4BP2L2 antibody.



Immunohistochemical analysis of paraffin-embedded human oophoroma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).