

Olfactory receptor 8K3 Polyclonal Antibody

Catalog No :	YT3451
Reactivity :	Human
Applications :	WB;IF;ELISA
Target :	Olfactory receptor 8K3
Fields :	>>Olfactory transduction
Gene Name :	OR8K3
Protein Name :	Olfactory receptor 8K3
Human Gene Id :	219473
Human Swiss Prot	Q8NH51
Immunogen :	The antiserum was produced against synthesized peptide derived from human OR8K3. AA range:236-285
Specificity :	Olfactory receptor 8K3 Polyclonal Antibody detects endogenous levels of Olfactory receptor 8K3 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)



Best Tools for immunology Research

Observed Band : 33kD

Cell Pathway : Olfactory transduction;

Background : Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. This olfactory receptor gene is a segregating pseudogene, where some individuals have an allele that encodes a functional olfactory receptor, while other individuals have an allele encoding a

 Function :
 function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,

Cell membrane; Multi-pass membrane protein.

Subcellular Location :

Products Images



Western Blot analysis of LOVO cells using Olfactory receptor 8K3 Polyclonal Antibody





Immunofluorescence analysis of HUVEC cells, using OR8K3 Antibody. The picture on the right is blocked with the synthesized peptide.

Western blot analysis of lysates from LOVO cells, using OR8K3 Antibody. The lane on the right is blocked with the synthesized peptide.