

Rho G Polyclonal Antibody

Catalog No :	YT4080
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	Rho G
Fields :	>>Bacterial invasion of epithelial cells;>>Salmonella infection;>>Yersinia infection
Gene Name :	RHOG
Protein Name :	Rho-related GTP-binding protein RhoG
Human Gene Id :	391
Human Swiss Prot No :	P84095
Mouse Gene Id :	56212
Mouse Swiss Prot No :	P84096
Immunogen :	The antiserum was produced against synthesized peptide derived from human RHOG. AA range:97-146
Specificity :	Rho G Polyclonal Antibody detects endogenous levels of Rho G protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 23kD


Background : This gene encodes a member of the Rho family of small GTPases, which cycle between inactive GDP-bound and active GTP-bound states and function as molecular switches in signal transduction cascades. Rho proteins promote reorganization of the actin cytoskeleton and regulate cell shape, attachment, and motility. The encoded protein facilitates translocation of a functional guanine nucleotide exchange factor (GEF) complex from the cytoplasm to the plasma membrane where ras-related C3 botulinum toxin substrate 1 is activated to promote lamellipodium formation and cell migration. Two related pseudogene have been identified on chromosomes 20 and X. [provided by RefSeq, Aug 2011],


Function : function:Required for the formation of membrane ruffles during macropinocytosis. Required for the formation of cup-like structures during trans-endothelial migration of leukocytes. In case of Salmonella enterica infection, activated by SopB and SGEF, which induces cytoskeleton rearrangements and promotes bacterial entry.,similarity:Belongs to the small GTPase superfamily. Rho family.,subunit:Interacts with SGEF.,

Subcellular Location : Cell membrane ; Lipid-anchor ; Cytoplasmic side .

Expression : Brain,PCR rescued clones,

Products Images





Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4 ° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).