

## Rabaptin-5a Polyclonal Antibody

YT3947 **Catalog No:** 

Human; Mouse; Rat Reactivity:

**Applications:** IHC;IF;ELISA

Target: Rabaptin-5a

Fields: >>Endocytosis

**Gene Name:** RABEP1

**Protein Name:** Rab GTPase-binding effector protein 1

Q15276

O35551

**Human Gene Id:** 9135

**Human Swiss Prot** 

No:

Mouse Gene Id: 54189

**Mouse Swiss Prot** 

No:

Rat Gene Id: 54190

Rat Swiss Prot No: 035550

Immunogen: The antiserum was produced against synthesized peptide derived from human

RABEP1. AA range:549-598

**Specificity:** Rabaptin-5a Polyclonal Antibody detects endogenous levels of Rabaptin-5a

protein.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200

1/3



**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 99kD

**Cell Pathway:** Endocytosis;

**Background:** function:Rab effector protein acting as linker between gamma-adaptin, RAB4A

and RAB5A. Involved in endocytic membrane fusion and membrane trafficking of recycling endosomes. Stimulates RABGEF1 mediated nucleotide exchange on RAB5A.,PTM:Proteolytic cleavage by caspases in apoptotic cells causes loss of

endosome fusion activity., similarity: Belongs to the rabaptin

family., subunit: Heterodimer with RABGEF1. The heterodimer binds RAB4A and RAB5A that have been activated by GTP-binding. Binds TSC2, GGA1, GGA2,

GGA3, AP1G1 and AP1G2.,

**Function:** function:Rab effector protein acting as linker between gamma-adaptin, RAB4A

and RAB5A. Involved in endocytic membrane fusion and membrane trafficking of recycling endosomes. Stimulates RABGEF1 mediated nucleotide exchange on RAB5A.,PTM:Proteolytic cleavage by caspases in apoptotic cells causes loss of

endosome fusion activity., similarity: Belongs to the rabaptin

family.,subunit:Heterodimer with RABGEF1. The heterodimer binds RAB4A and RAB5A that have been activated by GTP-binding. Binds TSC2, GGA1, GGA2,

GGA3, AP1G1 and AP1G2.,


Subcellular Location:

Cytoplasm. Early endosome. Recycling endosome. Cytoplasmic vesicle.

**Expression:** 

Cervix carcinoma, Epithelium, Fetal brain cortex, Kidney, Plate

## **Products Images**



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using RABEP1 Antibody. The picture on the right is blocked with the synthesized peptide.