

Synapsin I Polyclonal Antibody

Catalog No: YT4482

Reactivity: Human; Mouse; Rat

Applications: WB;IHC;IF;ELISA

Target: Synapsin I

Gene Name: SYN1

Protein Name: Synapsin-1

P17600

O88935

Human Gene Id: 6853

Human Swiss Prot

No:

Mouse Gene Id: 20964

Mouse Swiss Prot

No:

Rat Gene ld: 24949

Rat Swiss Prot No: P09951

Immunogen: The antiserum was produced against synthesized peptide derived from human

Synapsin1. AA range:26-75

Specificity: Synapsin I Polyclonal Antibody detects endogenous levels of Synapsin I protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not

yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 80kD

Background: This gene is a member of the synapsin gene family. Synapsins encode neuronal

phosphoproteins which associate with the cytoplasmic surface of synaptic vesicles. Family members are characterized by common protein domains, and they are implicated in synaptogenesis and the modulation of neurotransmitter release, suggesting a potential role in several neuropsychiatric diseases. This member of the synapsin family plays a role in regulation of axonogenesis and synaptogenesis. The protein encoded serves as a substrate for several different protein kinases and phosphorylation may function in the regulation of this protein in the nerve terminal. Mutations in this gene may be associated with X-linked disorders with primary neuronal degeneration such as Rett syndrome. Alternatively spliced transcript variants encoding different isoforms have been

identified. [provided by RefSeq, Jul 2008],

Function: disease:Defects in SYN1 are a cause of epilepsy X-linked with variable learning

disabilities and behavior disorders [MIM:300491]. XELBD is characterized by variable combinations of epilepsy, learning difficulties, macrocephaly, and aggressive behavior.,function:Neuronal phosphoprotein that coats synaptic vesicles, binds to the cytoskeleton, and is believed to function in the regulation of neurotransmitter release. The complex formed with NOS1 and CAPON proteins

is necessary for specific nitric-oxid functions at a presynaptic

level.,PTM:Substrate of at least four different protein kinases. It is probable that phosphorylation plays a role in the regulation of synapsin-1 in the nerve terminal. Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the synapsin family.,subunit:Homodimer. Interacts with CAPON. Forms a

ternary complex with NOS1. Isoform Ib interacts with


Subcellular Location:

Cell junction, synapse. Golgi apparatus.


Expression:

Brain, Brain cortex,


Products Images



Western Blot analysis of NIH-3T3 cells using Synapsin I Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Synapsin1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from NIH/3T3 cells, treated with Nocodazole 1ug/ml 16h, using Synapsin1 Antibody. The lane on the right is blocked with the synthesized peptide.