


## Ribosomal Protein L12 Polyclonal Antibody


Catalog No :	YT4095
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	Ribosomal Protein L12
Fields :	>>Ribosome;>>Coronavirus disease - COVID-19
Gene Name :	RPL12
Protein Name :	60S ribosomal protein L12
Human Gene Id :	6136
Human Swiss Prot	P30050
No : Mouse Gene Id :	269261
Mouse Swiss Prot	P35979
No : Rat Gene Id :	499782
Rat Swiss Prot No :	P23358
Immunogen :	The antiserum was produced against synthesized peptide derived from human RPL12. AA range:61-110
Specificity :	Ribosomal Protein L12 Polyclonal Antibody detects endogenous levels of Ribosomal Protein L12 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000 IF 1:50-200



Purification :	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	18kD
Cell Pathway :	Ribosome;
Background :	Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L11P family of ribosomal proteins. It is located in the cytoplasm. The protein binds directly to the 26S rRNA. This gene is co-transcribed with the U65 snoRNA, which is located in its fourth intron. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Jul 2008],
Function :	function:Binds directly to 26S ribosomal RNA.,similarity:Belongs to the ribosomal protein L11P family.,
Subcellular	nucleolus,cytosol,focal adhesion,membrane,cytosolic large ribosomal
Location :	subunit,extracellular matrix,extracellular exosome,
Expression :	Brain,Epithelium,Ovary,








Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.





Western blot analysis of lysates from HepG2 and HeLa cells, using RPL12 Antibody. The lane on the right is blocked with the synthesized peptide.