

SLC6A8 Polyclonal Antibody

Catalog No :	YT4318
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC
Target :	SLC6A8
Gene Name :	SLC6A8
Protein Name :	Sodium- and chloride-dependent creatine transporter 1
Human Gene Id :	6535
Human Swiss Prot No :	P48029
Mouse Gene Id :	102857
Mouse Swiss Prot No :	Q8VBW1
Rat Gene Id :	50690
Rat Swiss Prot No :	P28570
Immunogen :	The antiserum was produced against synthesized peptide derived from human SLC6A8. AA range:581-630
Specificity :	SLC6A8 Polyclonal Antibody detects endogenous levels of SLC6A8 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000;IHC 1:50-300
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 70kD


Background : The protein encoded by this gene is a plasma membrane protein whose function is to transport creatine into and out of cells. Defects in this gene can result in X-linked creatine deficiency syndrome. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2008],

Function : disease:Defects in SLC6A8 are the cause of X-linked creatine deficiency syndrome [MIM:300352]. X-linked creatine deficiency syndrome causes developmental delay, hypotonia, mental retardation, seizures, short stature and midface hypoplasia.,function:Required for the uptake of creatine in muscles and brain.,similarity:Belongs to the sodium:neurotransmitter symporter (SNF) family.,tissue specificity:Predominantly expressed in skeletal muscle and kidney. Also found in brain, heart, colon, testis and prostate.,


Subcellular Location : Membrane; Multi-pass membrane protein.

Expression : Predominantly expressed in skeletal muscle and kidney. Also found in brain, heart, colon, testis and prostate.


Products Images




Western Blot analysis of various cells using SLC6A8 Polyclonal Antibody



Western blot analysis of the lysates from HUVECcells using SLC6A8 antibody.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).