

Synaptotagmin XI Polyclonal Antibody

Catalog No :	YT4487
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	Synaptotagmin XI
Gene Name :	SYT11
Protein Name :	Synaptotagmin-11
Human Gene Id :	23208
Human Swiss Prot No :	Q9BT88
Mouse Gene Id :	229521
Mouse Swiss Prot No :	Q9R0N3
Rat Swiss Prot No :	O08835
Immunogen :	The antiserum was produced against synthesized peptide derived from human SYT11. AA range:181-230
Specificity :	Synaptotagmin XI Polyclonal Antibody detects endogenous levels of Synaptotagmin XI protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability : -15°C to -25°C/1 year (Do not lower than -25°C)

Observed Band : 48kD


Background : synaptotagmin 11 (SYT11) Homo sapiens This gene is a member of the synaptotagmin gene family and encodes a protein similar to other family members that are known calcium sensors and mediate calcium-dependent regulation of membrane trafficking in synaptic transmission. The encoded protein is also a substrate for ubiquitin-E3-ligase parkin. The gene has previously been referred to as synaptotagmin XII but has been renamed synaptotagmin XI to be consistent with mouse and rat official nomenclature. [provided by RefSeq, Apr 2010],

Function : cofactor: Binds 3 calcium ions per subunit. The ions are bound to the C2 domains., function: May be involved in Ca(2+)-dependent exocytosis of secretory vesicles through Ca(2+) and phospholipid binding to the C2 domain or may serve as Ca(2+) sensors in the process of vesicular trafficking and exocytosis., PTM: Ubiquitinated and targeted to the proteasome complex for degradation., similarity: Belongs to the synaptotagmin family., similarity: Contains 2 C2 domains., subcellular location: In substantia nigra, observed in neuronal cell bodies and neurites. Found in the core of the Lewy bodies in the brain of sporadic Parkinson disease patients., subunit: Homodimer. Can also form heterodimers (By similarity). Interacts with PARK2.,


Subcellular Location : Cytoplasmic vesicle membrane ; Single-pass membrane protein . Perikaryon . Golgi apparatus, trans-Golgi network membrane ; Single-pass membrane protein . Recycling endosome membrane ; Single-pass membrane protein . Lysosome membrane ; Single-pass membrane protein . Cytoplasmic vesicle, phagosome . Cell projection, axon . Cell projection, dendrite . Cell junction, synapse, postsynaptic density . Recycling endosome membrane ; Single-pass membrane protein . Cytoplasmic vesicle, clathrin-coated vesicle membrane ; Single-pass membrane protein . Perikaryon . Localized in vesicles that travels in axonal and dendritic shafts in both anterograde and retrograde directions. In macrophages and microglia, recruited in phagosomes at early stages of phagocytosis (By similarity). Found in the core of the

Expression : Amygdala, Bone marrow, Brain, Lymph,

Products Images



Western Blot analysis of various cells using Synaptotagmin XI Polyclonal Antibody



Western blot analysis of SYT11 Antibody. The lane on the right is blocked with the SYT11 peptide.