

## ZP4 Polyclonal Antibody

<b>Catalog No :</b>	YT4996
<b>Reactivity :</b>	Human;Rat;Mouse;
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	ZP4
<b>Gene Name :</b>	ZP4
<b>Protein Name :</b>	Zona pellucida sperm-binding protein 4
<b>Human Gene Id :</b>	57829
<b>Human Swiss Prot No :</b>	Q12836
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human ZP4. AA range:231-280
<b>Specificity :</b>	ZP4 Polyclonal Antibody detects endogenous levels of ZP4 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	65kD
<b>Background :</b>	The zona pellucida is an extracellular matrix that surrounds the oocyte and early embryo. It is composed primarily of three or four glycoproteins with various

functions during fertilization and preimplantation development. The nascent protein contains a N-terminal signal peptide sequence, a conserved ZP domain, a consensus furin cleavage site, and a C-terminal transmembrane domain. It is hypothesized that furin cleavage results in release of the mature protein from the plasma membrane for subsequent incorporation into the zona pellucida matrix. However, the requirement for furin cleavage in this process remains controversial based on mouse studies. Previously, this gene has been referred to as ZP1 or ZPB and thought to have similar functions as mouse Zp1. However, a human gene with higher similarity and chromosomal synteny to mouse Zp1 has been assigned the symbol ZP1 and this gene has been

## Function :

domain:The ZP domain is involved in the polymerization of the ZP proteins to form the zona pellucida.,function:The mammalian zona pellucida, which mediates species-specific sperm binding, induction of the acrosome reaction and prevents post-fertilization polyspermy, is composed of three to four glycoproteins, ZP1, ZP2, ZP3, and ZP4. ZP4 may act as a sperm receptor.,PTM:Proteolytically cleaved before the transmembrane segment to yield the secreted ectodomain incorporated in the zona pellucida.,similarity:Belongs to the ZP domain family. ZPB subfamily.,similarity:Contains 1 P-type (trefoil) domain.,similarity:Contains 1 ZP domain.,tissue specificity:Oocytes.,


## Subcellular Location :


[Processed zona pellucida sperm-binding protein 4]: Zona pellucida .; Cell membrane ; Single-pass type I membrane protein .

## Expression :

Expressed in oocytes.

## Products Images





Western blot analysis of the lysates from HepG2 cells using ZP4 antibody.