

TNF- β Polyclonal Antibody

Catalog No :	YT4691
Reactivity :	Human;Mouse;Rat;Monkey
Applications :	WB;ELISA
Target :	TNFB
Fields :	>>Cytokine-cytokine receptor interaction;>>Viral protein interaction with cytokine and cytokine receptor;>>NF-kappa B signaling pathway;>>TNF signaling pathway;>>Type I diabetes mellitus;>>Human T-cell leukemia virus 1 infection;>>Herpes simplex virus 1 infection
Gene Name :	LTA
Protein Name :	Lymphotoxin-alpha
Human Gene Id :	4049
Human Swiss Prot No :	P01374
Mouse Gene Id :	16992
Mouse Swiss Prot No :	P09225
Rat Gene Id :	25008
Rat Swiss Prot No :	Q06332
Immunogen :	The antiserum was produced against synthesized peptide derived from human TNF-beta. AA range:17-66
Specificity :	TNF- β Polyclonal Antibody detects endogenous levels of TNF- β protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG

Dilution : WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications.

Purification : The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 22kD

Cell Pathway : Cytokine-cytokine receptor interaction;Antigen processing and presentation;Type I diabetes mellitus;


Background : The encoded protein, a member of the tumor necrosis factor family, is a cytokine produced by lymphocytes. The protein is highly inducible, secreted, and forms heterotrimers with lymphotoxin-beta which anchor lymphotoxin-alpha to the cell surface. This protein also mediates a large variety of inflammatory, immunostimulatory, and antiviral responses, is involved in the formation of secondary lymphoid organs during development and plays a role in apoptosis. Genetic variations in this gene are associated with susceptibility to leprosy type 4, myocardial infarction, non-Hodgkin's lymphoma, and psoriatic arthritis. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, Jul 2012],

Function : disease:Genetic variations in LTA are associated with susceptibility to leprosy type 4 (LPRS4) [MIM:610988]; also known as susceptibility to age-dependent leprosy. Leprosy is a chronic infectious disease of peripheral sensory nerves caused by Mycobacterium leprae.,disease:Genetic variations in LTA are associated with susceptibility to psoriatic arthritis [MIM:607507]. Psoriasis is a chronic inflammatory dermatosis that affects approximately 2% of the population. It is characterized by red, scaly skin lesions that are usually found on the scalp, elbows, and knees, and may be associated with severe arthritis. Psoriatic arthritis has been defined as an inflammatory arthritis usually without any rheumatoid factor in serum (seronegative arthritis) associated with psoriasis.,function:Cytokine that in its homotrimeric form binds to TNFRSF1A/TNFR1, TNFRSF1B/TNFR and TNFRSF14/HVEM. In its hetero


Subcellular Location : Secreted. Membrane. The homotrimer is secreted. The heterotrimer is membrane-associated.

Expression : Lymph,Peripheral blood leukocyte,Peripheral blood mononuclear cells,Peripheral blood mononu

Products Images



Western Blot analysis of various cells using TNF- β Polyclonal Antibody diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysate from COS-7cell, using TNF- β antibodies.