

## **TP53INP2 Polyclonal Antibody**

Catalog No: YT4706

Reactivity: Human; Mouse; Rat

**Applications:** WB;IHC;IF;ELISA

Target: TP53INP2

Fields: >>Autophagy - animal

Gene Name: TP53INP2

**Protein Name:** Tumor protein p53-inducible nuclear protein 2

Q8IXH6

Q8CFU8

Human Gene Id: 58476

**Human Swiss Prot** 

No:

Mouse Gene ld: 68728

**Mouse Swiss Prot** 

No:

Rat Swiss Prot No: Q8CHM3

**Immunogen:** The antiserum was produced against synthesized peptide derived from human

TP53INP2. AA range:1-50

**Specificity:** TP53INP2 Polyclonal Antibody detects endogenous levels of TP53INP2 protein.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500-2000 IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet

tested in other applications.

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



**Concentration**: 1 mg/ml

**Storage Stability:** -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 24kD

Observed Band: 17kD

**Background:** tumor protein p53 inducible nuclear protein 2(TP53INP2) Homo sapiens The

protein encoded by this gene promotes autophagy and is essential for proper autophagosome formation and processing. In addition, the encoded protein can enhance rDNA transcription by helping in the assembly of the POLR1/RNA polymerase I preinitiation complex. Finally, this protein serves as a transcriptional

activator for some genes. [provided by RefSeq, Jul 2016],


Subcellular Location:

Cytoplasm, cytosol. Nucleus. Nucleus, PML body. Cytoplasmic vesicle, autophagosome. Shuttles between the nucleus and the cytoplasm, depending on cellular stress conditions, and re-localizes to autophagosomes on autophagy


activation.

**Expression :** Eye, Heart,

## **Products Images**



Immunofluorescence analysis of MCF7 cells, using TP53INP2 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using TP53INP2 Antibody. The picture on the right is blocked with the synthesized peptide.