

VIP Polyclonal Antibody

Catalog No :	YT4883
Reactivity :	Human;Rat;Mouse;
Applications :	IHC;IF;WB;ELISA
Target :	VIP
Fields :	>>cAMP signaling pathway;>>Neuroactive ligand-receptor interaction
Gene Name :	VIP
Protein Name :	VIP peptides
Human Gene Id :	7432
Human Swiss Prot No :	P01282
Mouse Swiss Prot No :	P32648
Immunogen :	The antiserum was produced against synthesized peptide derived from human VIP. AA range:71-120
Specificity :	VIP Polyclonal Antibody detects endogenous levels of VIP protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000 IHC 1:100 - 1:300. ELISA: 1:10000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 20kD


Background : The protein encoded by this gene belongs to the glucagon family. It stimulates myocardial contractility, causes vasodilation, increases glycogenolysis, lowers arterial blood pressure and relaxes the smooth muscle of trachea, stomach and gall bladder. The protein also acts as an antimicrobial peptide with antibacterial and antifungal activity. Alternative splicing occurs at this locus and two transcript variants encoding distinct isoforms have been identified. [provided by RefSeq, Nov 2014],

Function : function:PHM and PHV also cause vasodilation.,function:VIP causes vasodilation, lowers arterial blood pressure, stimulates myocardial contractility, increases glycogenolysis and relaxes the smooth muscle of trachea, stomach and gall bladder.,online information:Vasoactive intestinal peptide entry,similarity:Belongs to the glucagon family.,


Subcellular Location : Secreted.

Expression : Pancreatic carcinoma,Pheochromocytoma,Prostate,


Products Images



Immunohistochemical analysis of paraffin-embedded Human testis. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.



Immunohistochemistry analysis of VIP antibody in paraffin-embedded human testis tissue.



Western blot analysis of lysates from PC12 cells, primary antibody was diluted at 1:1000, 4° over night