

Btk Polyclonal Antibody

Catalog No :	YT5346
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	Btk
Fields :	>>NF-kappa B signaling pathway;>>Osteoclast differentiation;>>Platelet activation;>>B cell receptor signaling pathway;>>Fc epsilon RI signaling pathway;>>Epstein-Barr virus infection;>>Primary immunodeficiency
Gene Name :	BTK
Protein Name :	Tyrosine-protein kinase BTK
Human Gene Id :	695
Human Swiss Prot No :	Q06187
Mouse Gene Id :	12229
Mouse Swiss Prot No :	P35991
Immunogen :	Synthesized peptide derived from Btk . at AA range: 490-570
Specificity :	Btk Polyclonal Antibody detects endogenous levels of Btk protein only when non-phosphorylation at Y550.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)**Observed Band :** 76kD**Cell Pathway :** B_Cell_Antigen;Fc epsilon RI;Primary immunodeficiency;**Background :**

The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this gene cause X-linked agammaglobulinemia type 1, which is an immunodeficiency characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2013],

Function :

catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,cofactor:Binds 1 zinc ion per subunit.,disease:Defects in BTK are the cause of X-linked agammaglobulinemia (XLA) [MIM:300755]; also called X-linked agammaglobulinemia type 1 (AGMX1) or immunodeficiency type 1 (IMD1). XLA is a humoral immunodeficiency disease which results in developmental defects in the maturation pathway of B-cells. Affected boys have normal levels of pre-B-cells in their bone marrow but virtually no circulating mature B-lymphocytes. This results in a lack of immunoglobulins of all classes and leads to recurrent bacterial infections like otitis, conjunctivitis, dermatitis, sinusitis in the first few years of life, or even some patients present overwhelming sepsis or meningitis, resulting in death in a few hours. Treatment in most cases is by infusion of intravenous immunoglobulin.,



Subcellular Location :

Cytoplasm. Cell membrane; Peripheral membrane protein. Nucleus. In steady state, BTK is predominantly cytosolic. Following B-cell receptor (BCR) engagement by antigen, translocates to the plasma membrane through its PH domain. Plasma membrane localization is a critical step in the activation of BTK. A fraction of BTK also shuttles between the nucleus and the cytoplasm, and nuclear export is mediated by the nuclear export receptor CRM1.

Expression : Predominantly expressed in B-lymphocytes.

Products Images

Yang, Chunhui, et al. "Non-classical MHC IE negatively regulates macrophage activation and Th17 cell development in NOD mice." *Scientific reports* 5 (2015): 12941.



Western Blot analysis of 293 cells using Btk Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000