

### **BNIP-3 Polyclonal Antibody**

Catalog No: YT5154

**Reactivity:** Human; Mouse

**Applications:** WB;IHC;IF;ELISA

Target: BNIP-3

Fields: >>FoxO signaling pathway;>>Mitophagy - animal;>>Autophagy -

animal;>>Shigellosis;>>Legionellosis

Gene Name: BNIP3

**Protein Name:** BCL2/adenovirus E1B 19 kDa protein-interacting protein 3

Human Gene Id: 664

**Human Swiss Prot** 

No:

**Mouse Swiss Prot** 

No:

**Immunogen:** The antiserum was produced against synthesized peptide derived from the

Internal region of human BNIP3. AA range:71-120

**Specificity:** BNIP-3 Polyclonal Antibody detects endogenous levels of BNIP-3 protein.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Q12983

O55003

**Dilution :** WB 1:500 - 1:2000. IHC: 1:100-300 ELISA: 1:20000.. IF 1:50-200

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/3

**Observed Band:** 

22kD

### **Background:**

This gene is encodes a mitochondrial protein that contains a BH3 domain and acts as a pro-apoptotic factor. The encoded protein interacts with anti-apoptotic proteins, including the E1B 19 kDa protein and Bcl2. This gene is silenced in tumors by DNA methylation. [provided by RefSeq, Dec 2014],

#### **Function:**

function:Apoptosis-inducing protein that, which can overcome BCL2 suppression. May play a role in repartitioning calcium between the two major intracellular calcium stores in association with BCL2.,similarity:Belongs to the NIP3 family.,subcellular location:Coexpression with the EIB 19-kDa protein results in a shift in NIP3 localization pattern to the nuclear envelope.,subunit:Homodimer. Binds to BCL2. Interacts with BNIP3L. Also can

interact with adenovirus E1B 19 kDa protein or Epstein-Barr virus BHRF1.,


# Subcellular Location:

Mitochondrion. Mitochondrion outer membrane; Single-pass membrane protein. Coexpression with the EIB 19-kDa protein results in a shift in NIP3 localization pattern to the nuclear envelope. Colocalizes with ACAA2 in the mitochondria. Colocalizes with SPATA18 at the mitochondrion outer membrane.


### **Expression:**

B-cell, Brain, Cerebellum, Kidney, Testis,


## **Products Images**




Western Blot analysis of mouse eye cells using BNIP-3 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded humanbrain, antibody was diluted at 1:100



Western blot analysis of lysate from mouse eye cells, using BNIP3 Antibody.



Feng, Qin. "Quantitative proteomic analysis reveals that Arctigenin alleviates concanavalin A-induced hepatitis through suppressing immune system and regulating autophagy." Frontiers in immunology 9 (2018): 1881.

