

gp91-phox Polyclonal Antibody


Catalog No :	YT5383
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	gp91-phox
Fields :	>>HIF-1 signaling pathway;>>Phagosome;>>Ferroptosis;>>Necroptosis;>>Neutrophil extracellular trap formation;>>NOD-like receptor signaling pathway;>>Leukocyte transendothelial migration;>>AGE-RAGE signaling pathway in diabetic complications;>>Alzheimer disease;>>Prion disease;>>Pathways of neurodegeneration - multiple diseases;>>Leishmaniasis;>>Coronavirus disease - COVID-19;>>Diabetic cardiomyopathy;>>Lipid and atherosclerosis
Gene Name :	CYBB
Protein Name :	Cytochrome b-245 heavy chain
Human Gene Id :	1536
Human Swiss Prot No :	P04839
Mouse Swiss Prot No :	Q61093
Immunogen :	The antiserum was produced against synthesized peptide derived from the Internal region of human CYBB. AA range:111-160
Specificity :	gp91-phox Polyclonal Antibody detects endogenous levels of gp91-phox protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC: 1:100-1:300. ELISA: 1:20000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	70kD
Cell Pathway :	Leukocyte transendothelial migration;
Background :	<p>Cytochrome b (-245) is composed of cytochrome b alpha (CYBA) and beta (CYBB) chain. It has been proposed as a primary component of the microbicidal oxidase system of phagocytes. CYBB deficiency is one of five described biochemical defects associated with chronic granulomatous disease (CGD). In this disorder, there is decreased activity of phagocyte NADPH oxidase; neutrophils are able to phagocytize bacteria but cannot kill them in the phagocytic vacuoles. The cause of the killing defect is an inability to increase the cell's respiration and consequent failure to deliver activated oxygen into the phagocytic vacuole. [provided by RefSeq, Jul 2008],</p>
Function :	<p>cofactor:FAD.,disease:Defects in CYBB are a cause of chronic granulomatous disease X-linked (XCGD) [MIM:306400]. Chronic granulomatous disease is a genetically heterogeneous disorder characterized by the inability of neutrophils and phagocytes to kill microbes that they have ingested. Patients suffer from life-threatening bacterial/fungal infections.,function:Critical component of the membrane-bound oxidase of phagocytes that generates superoxide. It is the terminal component of a respiratory chain that transfers single electrons from cytoplasmic NADPH across the plasma membrane to molecular oxygen on the exterior. Also functions as a voltage-gated proton channel that mediates the H(+) currents of resting phagocytes. It participates in the regulation of cellular pH and is blocked by zinc.,online information:CYBB deficiency database,PTM:Glycosylated.,similarity:Contains 1 FAD-binding FR-t</p>
Subcellular Location :	Cell membrane; Multi-pass membrane protein. As unassembled monomer may localize to the endoplasmic reticulum. .
Expression :	Detected in neutrophils (at protein level).


Products Images



Western Blot analysis of K562 cells using gp91-phox Polyclonal Antibody. Antibody was diluted at 1:2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western Blot analysis of HEPG2 using gp91-phox Polyclonal Antibody diluted at 1:2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysate from K562 cells, using CYBB Antibody.