

CD47 Polyclonal Antibody

Catalog No :	YT5509
Reactivity :	Human;Rat;Mouse;
Applications :	WB;IHC;IF;ELISA
Target :	CD47
Fields :	>>ECM-receptor interaction
Gene Name :	CD47
Protein Name :	Leukocyte surface antigen CD47
Human Gene Id :	961
Human Swiss Prot No :	Q08722
Mouse Swiss Prot No :	Q61735
Immunogen :	The antiserum was produced against synthesized peptide derived from the Internal region of human CD47. AA range:61-110
Specificity :	CD47 Polyclonal Antibody detects endogenous levels of CD47 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC: 1:100-1:300. ELISA: 1:20000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 35kD

Cell Pathway : ECM-receptor interaction;


Background : This gene encodes a membrane protein, which is involved in the increase in intracellular calcium concentration that occurs upon cell adhesion to extracellular matrix. The encoded protein is also a receptor for the C-terminal cell binding domain of thrombospondin, and it may play a role in membrane transport and signal transduction. This gene has broad tissue distribution, and is reduced in expression on Rh erythrocytes. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2010],

Function : function:Has a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in the modulation of integrins. Plays an important role in memory formation and synaptic plasticity in the hippocampus (By similarity). Receptor for SIRPA, binding to which prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. Interaction with SIRPG mediates cell-cell adhesion, enhances superantigen-dependent T-cell-mediated proliferation and costimulates T-cell activation. May play a role in membrane transport and/or integrin dependent signal transduction. May prevent premature elimination of red blood cells. May be involved in membrane permeability changes induced following virus infection.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,subunit:Interacts with THBS1 and fibrinogen (By similarity). Monomer. Inter


Subcellular Location : Cell membrane ; Multi-pass membrane protein .

Expression : Very broadly distributed on normal adult tissues, as well as ovarian tumors, being especially abundant in some epithelia and the brain.

Products Images



Western Blot analysis of 3T3 cells using CD47 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-Lymph-nodes, antibody was diluted at 1:100