

## TNFAIP8L3 Polyclonal Antibody

<b>Catalog No :</b>	YT6072
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	TNFAIP8L3
<b>Gene Name :</b>	TNFAIP8L3
<b>Protein Name :</b>	TNFAIP8L3
<b>Human Gene Id :</b>	388121
<b>Human Swiss Prot No :</b>	Q5GJ75
<b>Mouse Gene Id :</b>	244882
<b>Mouse Swiss Prot No :</b>	Q3TBL6
<b>Immunogen :</b>	Synthesized peptide derived from human TNFAIP8L3. at AA range: 15-45
<b>Specificity :</b>	This antibody detects endogenous levels of TNFAIP8L3
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000, ELISA 1:10000-20000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

**Observed Band :** 33kD


**Background :** caution:It is uncertain whether Met-1 or Met-89 is the initiator.,similarity:Belongs to the TNFAIP8 family.,

**Function :** caution:It is uncertain whether Met-1 or Met-89 is the initiator.,similarity:Belongs to the TNFAIP8 family.,

**Subcellular Location :** Cytoplasm . Cell membrane . On PDGF activation, translocates from cytoplasm to plasma membrane. .

**Expression :** Widely expressed (at protein level) (PubMed:25479791). Highly expressed in most carcinoma cell lines (PubMed:25479791, PubMed:25242044).

## Products Images



Western Blot analysis of 1,293T 2,mouse-brain cells using primary antibody diluted at 1:1000(4 °C overnight). Secondary antibody:Goat Anti-rabbit IgG IRDye 800( diluted at 1:5000, 25 °C, 1 hour)