

PARP-1 Polyclonal Antibody

Catalog No :	YT6210
Reactivity :	Human;Mouse;Rat
Applications :	IHC;IF;WB
Target :	PARP
Fields :	>>Base excision repair;>>NF-kappa B signaling pathway;>>Apoptosis;>>Necroptosis;>>Diabetic cardiomyopathy
Gene Name :	PARP1 ADPRT PPOL
Protein Name :	PARP-1
Human Gene Id :	142
Human Swiss Prot No :	P09874
Immunogen :	Synthesized peptide derived from human PARP-1. AA range: 410-460
Specificity :	This antibody detects endogenous levels of human PARP-1
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	IHC 1:50-200, WB 1:500-2000. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	113kD


Background : This gene encodes a chromatin-associated enzyme, poly(ADP-riboseyl)transferase, which modifies various nuclear proteins by poly(ADP-riboseyl)ation. The modification is dependent on DNA and is involved in the regulation of various important cellular processes such as differentiation, proliferation, and tumor transformation and also in the regulation of the molecular events involved in the recovery of cell from DNA damage. In addition, this enzyme may be the site of mutation in Fanconi anemia, and may participate in the pathophysiology of type I diabetes. [provided by RefSeq, Jul 2008],

Function : catalytic activity:NAD(+) + (ADP-D-riboseyl)(n)-acceptor = nicotinamide + (ADP-D-riboseyl)(n+1)-acceptor.,function:Involved in the base excision repair (BER) pathway, by catalyzing the poly(ADP-riboseyl)ation of a limited number of acceptor proteins involved in chromatin architecture and in DNA metabolism. This modification follows DNA damages and appears as an obligatory step in a detection/signaling pathway leading to the reparation of DNA strand breaks.,miscellaneous:The ADP-D-riboseyl group of NAD(+) is transferred to an acceptor carboxyl group on a histone or the enzyme itself, and further ADP-riboseyl groups are transferred to the 2'-position of the terminal adenosine moiety, building up a polymer with an average chain length of 20-30 units.,PTM:Phosphorylated by PRKDC. Phosphorylated upon DNA damage, probably by ATM or ATR.,PTM:Poly-ADP-riboseylated by PARP2.,similarity:Contains 1 BRCT


Subcellular Location : Nucleus . Nucleus, nucleolus . Chromosome . Localizes to sites of DNA damage.

Expression : Brain,Colon carcinoma,Fibroblast,Lung,Ovarian carcinoma,Skin,


Products Images




Zhou, Li, et al. "Coptisine induces apoptosis in human hepatoma cells through activating 67-kDa laminin receptor/cGMP signaling." *Frontiers in pharmacology* 9 (2018).



Yao, Chong, et al. "Crocin induces autophagic apoptosis in hepatocellular carcinoma by inhibiting Akt/mTOR activity." *OncoTargets and therapy* 11 (2018): 2017.



Zhou, Yanlong, et al. "The degradation of mixed lineage kinase domain-like protein promotes neuroprotection after ischemic brain injury." *Oncotarget* 8.40 (2017): 68393.



Immunohistochemical analysis of paraffin-embedded human brain. 1, Tris-EDTA, pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200(4° overnight).3,Secondary antibody was diluted at 1:200(room temperature, 45min).