


## COL14A1 Polyclonal Antibody

Catalog No :	YT5999
Reactivity :	Human;Mouse;Rat
Applications :	IHC;IF;ELISA
Target :	COL14A1
Fields :	>>Protein digestion and absorption
Gene Name :	COL14A1 UND
Protein Name :	Collagen alpha-1(XIV) chain (Undulin)
Human Gene Id :	7373
fruinan Generu .	1010
Human Swiss Prot No :	Q05707
Mouse Swiss Prot	Q80X19
Immunogen :	Synthetic peptide from human c-ternal protein at AA range: 1571-1620
Specificity :	The antibody detects endogenous COL14A1
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	IHC 1:50-200, ELISA 1:10000-20000. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)



Background :	This gene encodes the alpha chain of type XIV collagen, a member of the FACIT (fibril-associated collagens with interrupted triple helices) collagen family. Type XIV collagen interacts with the fibril surface and is involved in the regulation of fibrillogenesis. [provided by RefSeq, Jan 2013],
Function :	function:Plays an adhesive role by integrating collagen bundles. It is probably associated with the surface of interstitial collagen fibrils via COL1. The COL2 domain may then serve as a rigid arm which sticks out from the fibril and protrudes the large N-terminal globular domain into the extracellular space, where it might interact with other matrix molecules or cell surface receptors.,PTM:Lysines at the third position of the tripeptide repeating unit (G-X- Y) are hydroxylated in all cases and bind carbohydrates.,PTM:May contain numerous cysteine residues involved in inter- and intramolecular disulfide bonding.,PTM:Prolines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains.,similarity:Belongs to the fibril- associated collagens with interrupted helices (FACIT) family.,similarity:Contains 1 TSP N-terminal (TSPN) domain.,similarity:
Subcellular Location :	Secreted, extracellular space, extracellular matrix .
Expression :	Brain,Liver,Muscle,PNS,Testis,

## Products Images



Immunohistochemical analysis of paraffin-embedded humanskin, antibody was diluted at 1:200