

TPM4 rabbit pAb

Catalog No :	YT6301
Reactivity :	Human;Mouse;Rat
Applications :	WB
Target :	TPM4
Fields :	>>Cardiac muscle contraction;>>Adrenergic signaling in cardiomyocytes;>>Hypertrophic cardiomyopathy;>>Dilated cardiomyopathy
Gene Name :	TPM4
Protein Name :	TPM4
Human Gene Id :	7171
Human Swiss Prot No :	P67936
Mouse Gene Id :	326618
Mouse Swiss Prot No :	Q6IRU2
Rat Gene Id :	24852
Rat Swiss Prot No :	P09495
Immunogen :	Synthesized peptide derived from human TPM4 AA range: 127-177
Specificity :	This antibody detects endogenous levels of TPM4 at Human/Mouse/Rat
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000

Purification : The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 27kD


Background : This gene encodes a member of the tropomyosin family of actin-binding proteins involved in the contractile system of striated and smooth muscles and the cytoskeleton of non-muscle cells. Tropomyosins are dimers of coiled-coil proteins that polymerize end-to-end along the major groove in most actin filaments. They provide stability to the filaments and regulate access of other actin-binding proteins. In muscle cells, they regulate muscle contraction by controlling the binding of myosin heads to the actin filament. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2009],

Function : domain:The molecule is in a coiled coil structure that is formed by 2 polypeptide chains. The sequence exhibits a prominent seven-residues periodicity.,function:Binds to actin filaments in muscle and non-muscle cells. Plays a central role, in association with the troponin complex, in the calcium dependent regulation of vertebrate striated muscle contraction. Smooth muscle contraction is regulated by interaction with caldesmon. In non-muscle cells is implicated in stabilizing cytoskeleton actin filaments. Binds calcium.,similarity:Belongs to the tropomyosin family.,subunit:Heterodimer of an alpha and a beta chain.,tissue specificity:Detected in cardiac tissue and platelets, the form found in cardiac tissue is a higher molecular weight than the form found in platelets. Expressed at higher levels in the platelets of hypertensive patients with cardiac hypertrophy than in the platelets of hyp

Subcellular Location : Cytoplasm, cytoskeleton . Associates with F-actin stress fibers. .

Expression : Detected in cardiac tissue and platelets, the form found in cardiac tissue is a higher molecular weight than the form found in platelets. Expressed at higher levels in the platelets of hypertensive patients with cardiac hypertrophy than in the platelets of hypertensive patients without cardiac hypertrophy (at protein level).

Products Images



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night