

HCFC2 rabbit pAb

Catalog No: YT6455

Reactivity: Human; Mouse; Rat

Applications: WB

Target: HCFC2

Fields: >>Herpes simplex virus 1 infection

Q9Y5Z7

Q9D968

Gene Name: HCFC2

Protein Name: HCFC2

Human Gene Id: 29915

Human Swiss Prot

Tullian Swiss Fro

No:

Mouse Swiss Prot

No:

Rat Gene Id: 314704

Rat Swiss Prot No: Q5RKG2

Immunogen: Synthesized peptide derived from human HCFC2 AA range: 347-397

Specificity: This antibody detects endogenous levels of HCFC2 at Human/Mouse/Rat

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1 ? 500-2000

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 87kD

Background: This gene encodes one of two proteins which interact with VP16, a herpes

simplex virus protein that initiates virus infection. Both the encoded protein and the original Herpes host cell factor interact with VP16 through a beta-propeller domain. The original Herpes host cell factor, however, is effective at initiating viral infection while the encoded protein is not. Transcripts of varying length due to alternative polyadenylation signals have been described. [provided by RefSeq,

Jul 2008],

Function: similarity:Contains 3 fibronectin type-III domains.,similarity:Contains 4 Kelch

repeats., subunit: Binds MLL. Component of the MLL complex, at least composed of MLL, ASH2L, RBBP5, DPY30, WDR5, MEN1, HCFC1 and HCFC2., tissue specificity: Highly expressed in testis. Detected at lower levels in spleen, thymus,

prostate, ovary, small intestine and colon.,

Subcellular Location :

60 --45 --35 --25 --15 --10 -- Cytoplasm . Nucleus .

Expression: Highly expressed in testis. Detected at lower levels in spleen, thymus, prostate,

ovary, small intestine and colon.

Products Images

