

## TBL3 rabbit pAb

YT6754 **Catalog No:** 

Human; Mouse; Rat Reactivity:

**Applications:** WB

Target: TBL3

Fields: >>Ribosome biogenesis in eukaryotes

Q12788

Q8C4J7

Gene Name: TBL3 SAZD

**Protein Name:** TBL3

**Human Gene Id:** 10607

**Human Swiss Prot** 

No:

Mouse Gene Id: 213773

**Mouse Swiss Prot** 

No:

Rat Gene Id: 287120

Rat Swiss Prot No: Q5U2W5

Immunogen: Synthesized peptide derived from human TBL3 AA range: 486-536

This antibody detects endogenous levels of TBL3 at Human/Mouse/Rat **Specificity:** 

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1?500-2000

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

1/2



chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 89kD

**Background:** The protein encoded by this gene has sequence similarity with members of the

WD40 repeat-containing protein family. The WD40 group is a large family of proteins, which appear to have a regulatory function. It is believed that the WD40 repeats mediate protein-protein interactions and members of the family are involved in signal transduction, RNA processing, gene regulation, vesicular trafficking, cytoskeletal assembly and may play a role in the control of cytotypic differentiation. This gene has multiple polyadenylation sites. It might have multiple alternatively spliced transcript variants but the variants have not been fully


described yet. [provided by RefSeq, Jul 2008],

**Function:** similarity:Contains 13 WD repeats.,

Subcellular Location:

Nucleus, nucleolus.

## **Products Images**



Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4° over night