

PRDM2 rabbit pAb

Catalog No :	YT7018
Reactivity :	Human;Rat
Applications :	WB;IHC
Target :	PRDM2
Fields :	>>Lysine degradation;>>Metabolic pathways
Gene Name :	PRDM2 KMT8 RIZ
Protein Name :	PRDM2
Human Gene Id :	7799
Human Swiss Prot No :	Q13029
Rat Gene Id :	313678
Rat Swiss Prot No :	Q63755
Immunogen :	Synthesized peptide derived from human PRDM2 AA range: 964-1014
Specificity :	This antibody detects endogenous levels of PRDM2 at Human/Rat
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000;IHC 1:50-300
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 189kD

Background : This tumor suppressor gene is a member of a nuclear histone/protein methyltransferase superfamily. It encodes a zinc finger protein that can bind to retinoblastoma protein, estrogen receptor, and the TPA-responsive element (MTE) of the heme-oxygenase-1 gene. Although the functions of this protein have not been fully characterized, it may (1) play a role in transcriptional regulation during neuronal differentiation and pathogenesis of retinoblastoma, (2) act as a transcriptional activator of the heme-oxygenase-1 gene, and (3) be a specific effector of estrogen action. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2008],


Function : catalytic activity:S-adenosyl-L-methionine + histone L-lysine = S-adenosyl-L-homocysteine + histone N(6)-methyl-L-lysine.,function:S-adenosyl-L-methionine-dependent histone methyltransferase that specifically methylates 'Lys-9' of histone H3. May function as a DNA-binding transcription factor. Binds to the macrophage-specific TPA-responsive element (MTE) of the HMOX1 (heme oxygenase 1) gene and may act as a transcriptional activator of this gene.,similarity:Contains 1 SET domain.,similarity:Contains 8 C2H2-type zinc fingers.,subunit:Binds to the retinoblastoma protein (RB). Interacts with GATA3.,tissue specificity:Highly expressed in retinoblastoma cell lines and in brain tumors. Also expressed in a number of other cell lines and in brain, heart, skeletal muscle, liver and spleen. Isoform 1 is expressed in testis at much higher level than isoform 3.,

Subcellular Nucleus .


Location :

Expression : Highly expressed in retinoblastoma cell lines and in brain tumors. Also expressed in a number of other cell lines and in brain, heart, skeletal muscle, liver and spleen. Isoform 1 is expressed in testis at much higher level than isoform 3.

Products Images



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).