

ZN521 rabbit pAb

Catalog No :	YT7130
Reactivity :	Human;Mouse
Applications :	WB
Target :	ZN521
Gene Name :	ZNF521 EHZF LIP3
Protein Name :	ZN521
Human Gene Id :	25925
Human Swiss Prot No :	Q96K83
Mouse Gene Id :	225207
Mouse Swiss Prot No :	Q6KAS7
Immunogen :	Synthesized peptide derived from human ZN521 AA range: 997-1047
Specificity :	This antibody detects endogenous levels of ZN521 at Human/Mouse
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1[?]500-2000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 144kD**Function :**

disease:A chromosomal aberration involving ZNF521 is found in acute lymphoblastic leukemia. Translocation t(9;18)(p13;q11.2) with PAX5. The translocation generates the PAX5-ZNF521 oncogene consisting of the N-terminus part of PAX5 and the C-terminus part of ZNF521.,domain:Uses different DNA- and protein-binding zinc fingers to regulate the distinct BMP-Smad and hematopoietic system.,function:Transcription factor that can both act as an activator or a repressor depending on the context. Involved in BMP signaling and in the regulation of the immature compartment of the hematopoietic system. Associates with SMADs in response to BMP2 leading to activate transcription of BMP target genes. Acts as a transcriptional repressor via its interaction with EBF1, a transcription factor involved specification of B-cell lineage; this interaction preventing EBF1 to bind DNA and activate target genes.,sim


Subcellular Location :

Nucleus .

Expression :

Predominantly expressed in hematopoietic cells. Present in organs and tissues that contain stem and progenitor cells, myeloid and/or lymphoid: placenta, spleen, lymph nodes, thymus, bone marrow and fetal liver. Within the hematopoietic system, it is abundant in CD34(+) cells but undetectable in mature peripheral blood leukocytes, and its levels rapidly decrease during the differentiation of CD34(+) cells in response to hemopoietins.

Products Images



Western blot analysis of lysates from MDA-MB cells, primary antibody was diluted at 1:1000, 4° over night