

THRSP rabbit pAb

| | |
|------------------------------|---|
| Catalog No : | YT7277 |
| Reactivity : | Human;Mouse;Rat |
| Applications : | WB;IHC |
| Target : | THRSP |
| Gene Name : | THRSP |
| Protein Name : | THRSP |
| Human Gene Id : | 7069 |
| Human Swiss Prot No : | Q92748 |
| Mouse Gene Id : | 21835 |
| Mouse Swiss Prot No : | Q62264 |
| Rat Swiss Prot No : | P04143 |
| Immunogen : | Synthesized peptide derived from human THRSP AA range: 4-54 |
| Specificity : | This antibody detects endogenous levels of THRSP at Human/Mouse/Rat |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:500-2000;IHC 1:50-300 |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 16kD


Background : The protein encoded by this gene is similar to the gene product of S14, a rat gene whose expression is limited to liver and adipose tissue and is controlled by nutritional and hormonal factors. This gene has been shown to be expressed in liver and adipocytes, particularly in lipomatous modules. It is also found to be expressed in lipogenic breast cancers, which suggests a role in controlling tumor lipid metabolism. [provided by RefSeq, Jul 2008],

Function : function:May play an important role in lipogenesis.,similarity:Belongs to the SPOT14 family.,tissue specificity:Mainly expressed in tissues that synthesize triglycerides.,


Subcellular Location : Nucleus . Cytoplasm .

Expression : Mainly expressed in tissues that synthesize triglycerides.

Products Images



Western blot analysis of lysates from PC-12 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).