

## SLTM rabbit pAb


<b>Catalog No :</b>	YT7364
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB;IHC
<b>Target :</b>	SLTM
<b>Gene Name :</b>	SLTM MET
<b>Protein Name :</b>	SLTM
<b>Human Gene Id :</b>	79811
<b>Human Swiss Prot No :</b>	Q9NWH9
<b>Mouse Gene Id :</b>	66660
<b>Mouse Swiss Prot No :</b>	Q8CH25
<b>Immunogen :</b>	Synthesized peptide derived from human SLTM AA range: 562-612
<b>Specificity :</b>	This antibody detects endogenous levels of SLTM at Human/Mouse
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000;IHC 1:50-300
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

**Molecularweight :** 114kD


**Function :** function:When overexpressed, acts as a general inhibitor of transcription that eventually leads to apoptosis.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,sequence caution:Contaminating sequence. Potential poly-A sequence.,sequence caution:Translation N-terminally extended.,similarity:Contains 1 RRM (RNA recognition motif) domain.,similarity:Contains 1 SAP domain.,subcellular location:Detected in punctate structures.,

**Subcellular Location :** Nucleus . Detected in punctate structures. .

## Products Images



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).