

## CYBR1 rabbit pAb

Catalog No: YT7604

Reactivity: Human; Mouse; Rat

**Applications:** WB

Target: CYBR1

**Fields:** >>Mineral absorption

Gene Name: CYBRD1 DCYTB FRRS3

Q53TN4

Q925G2

Protein Name: CYBR1

Human Gene Id: 79901

**Human Swiss Prot** 

iuman Swiss Fio

No:

Mouse Gene Id: 73649

**Mouse Swiss Prot** 

No:

**Rat Gene Id:** 295669

Rat Swiss Prot No: Q5RKJ2

Immunogen: Synthesized peptide derived from human CYBR1 AA range: 41-91

**Specificity:** This antibody detects endogenous levels of CYBR1 at Human/Mouse/Rat

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1 ? 500-2000

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

1/3



chromatography using epitope-specific immunogen.

**Concentration:** 1 mg/ml

-15°C to -25°C/1 year(Do not lower than -25°C) **Storage Stability:** 

Molecularweight: 31kD

This gene is a member of the cytochrome b(561) family that encodes an iron-**Background:** 

> regulated protein. It highly expressed in the duodenal brush border membrane. It has ferric reductase activity and is believed to play a physiological role in dietary

iron absorption. [provided by RefSeq, Jul 2008],

**Function:** cofactor:Binds 2 heme groups non-covalently.,disease:Defects in CYBRD1 may

be a cause of primary hereditary hemochromatosis (HFE). HFE is an iron-loading

disorder characterized by iron accumulation in parenchymal cells. Iron

accumulation usually results in tissue damage and causes cirrhosis of the liver. diabetes mellitus, arthropathy, cardiomyopathy, endocrine abnormalities and an increased risk of hepatocellular carcinoma.,function:Ferric-chelate reductase that reduces Fe(3+) to Fe(2+). Present at the brush border of duodenal enterocytes where it probably reduces dietary Fe(3+) thereby facilitating its transport into the mucosal cells. Uses ascorbate as electron donor. May be involved in extracellular ascorbate recycling in erythrocyte membranes. May also act as a ferrireductase in

airway epithelial cells.,induction:By iron deficiency (at protein

level)., similarity: Contains 1 cytochrom

Subcellular

Cell membrane; Multi-pass membrane protein. Apical cell membrane; Multipass membrane protein. Localized at the brush border of duodenal cells... Location:


**Expression:** 

Present in erythrocyte membranes (at protein level). Also expressed in

respiratory epithelium.

## **Products Images**

2/3



Western blot analysis of lysates from 3T3 cells, primary antibody was diluted at 1:1000, 4° over night